

Memory T Cells Specific for Novel Human Papillomavirus Type 16 (HPV16) E6 Epitopes in Women Whose HPV16 Infection Has Become Undetectable[∇]

Xuelian Wang,^{1,2*} Anna-Barbara Moscicki,³ Laura Tsang,⁴
 Andrea Brockman,⁴ and Mayumi Nakagawa¹

Departments of Pathology¹ and Microbiology and Immunology,⁴ College of Medicine, University of Arkansas for Medical Sciences, Little Rock, Arkansas; Department of Microbiology and Parasitology, College of Basic Medical Sciences, China Medical University, Shenyang, China²; and Department of Pediatrics, College of Medicine, University of California at San Francisco, San Francisco, California³

Received 5 October 2007/Returned for modification 9 January 2008/Accepted 14 April 2008

Human papillomavirus (HPV)-specific T-cell response to the HPV type 16 (HPV16) E6 protein has been shown to be associated with successful viral clearance. The patterns of CD8 T-cell epitopes within HPV16 E6 protein were previously studied in two women with HPV16 clearance. The goal of this study was to characterize these epitopes in terms of their minimal and optimal amino acid sequences and the human leukocyte antigen (HLA) restriction molecules. The presence of the epitope-specific memory T cells after viral clearance was also examined. In subject A, the dominant epitope was characterized to be E6 75–83 (KFYSKISEY), restricted by the HLA-B62 molecule, while that of subject B was E6 133–142 (HNIRGRWTGR), restricted by the HLA-A6801 molecule. Homologous epitopes were identified in five other high-risk HPV types for both of these epitopes, but they were not recognized by respective T-cell clone cells. An enzyme-linked immunospot assay or tetramer analysis was performed on peripheral blood mononuclear cells from blood samples collected after viral clearance but prior to isolation of the T-cell clones. The presence of epitope-specific memory T cells was demonstrated. These data suggest that HPV-specific memory T cells were generated *in vivo* and that they may remain in circulation many months, if not years, after viral clearance. Our findings broaden the spectrum of the CD8 T-cell epitopes of the HPV16 E6 protein. The characterization of novel T-cell epitopes and long-lasting epitope-specific memory T cells may be useful for the development of a potential epitope-based vaccine.

Cervical cancer is the second most common malignant cancer in women worldwide. High-risk human papillomaviruses (HPVs), most commonly HPV type 16 (HPV16), have been shown as the major cause of cervical cancer (29, 33). The transforming properties of high-risk HPV16 E6 and E7 proteins play a critical role in the initiation of oncogenic processes. The E6 protein can lead to the degradation of the p53 tumor suppressor protein (4, 28), while the E7 protein can interfere with the retinoblastoma tumor suppressor protein (28). These two proteins are foreign viral antigens, and their constitutive expression is required for the maintenance of the malignant phenotype of cervical cancer (32, 33). Therefore, HPV16 E6 and E7 proteins have been sought as ideal targets of antigen-specific immunotherapeutic strategies for the prevention and treatment of cervical cancer.

Immune responses, especially HPV-specific T-cell immune responses, are thought to play an important role in successfully clearing HPV infection and controlling HPV-associated diseases. CD8-positive, major histocompatibility complex class I-restricted cytotoxic T lymphocytes (CTLs) are known to be responsible for recognizing and killing virus-infected host cells and virus-induced tumor cells (12). Immunohistochemical

analyses of squamous intraepithelial lesion and cervical cancer specimens have demonstrated the presence of activated CTLs in lesions (2). A study using stimulated peripheral blood mononuclear cells (PBMCs) from cervical cancer patients with a human leukocyte antigen A2 (HLA-A2)-restricted HPV16 E7 peptide (E7 11–20) showed that CTLs were capable of lysing HLA-matched HPV16 E7 11–20-pulsed targets in two of three patients (1). Evans et al. (6) identified HPV-specific CTLs in lymph nodes and tumors in cervical cancer patients. In our previous work examining CTL responses to HPV16 in HPV16-infected women (no squamous intraepithelial lesion), we showed that CTL responses to the HPV16 E6 protein, but not to the E7 protein, were significantly associated with the clearance of HPV16 infection (24). We also investigated the CD8 T-cell epitopes of HPV16 E6 from women in whom the presence of dominant HPV16 E6 CD8 T-cell epitopes were demonstrated to be in the E6 16–40 region (21, 22). A dominant region was defined as the region within the E6 protein in which the largest number of spot-forming units was demonstrated for a particular subject using the gamma interferon (IFN- γ) enzyme-linked immunospot (ELISPOT) assay (22). The goals of this study were to characterize the CD8 T-cell epitopes from subjects in whom dominant responses were shown to be in other E6 regions (22) and to demonstrate the presence of epitope-specific memory T cells. Two novel HPV16 E6 CD8 T-cell epitopes, E6 75–83 (KFYSKISEY), restricted by HLA-B62, and E6 133–142 (HNIRGRWTGR), restricted by HLA-A68, were characterized. The presence of HPV-specific mem-

* Corresponding author. Mailing address: Department of Pathology, University of Arkansas for Medical Sciences, 4301 West Markham Street, Slot 824, Little Rock, AR 72205. Phone: (501) 526-7812. Fax: (501) 526-5934. E-mail: xwang4@uams.edu.

[∇] Published ahead of print on 30 April 2008.

ory T cells was examined using an ELISPOT assay or tetramer staining, and the HPV-specific memory T cells appeared to remain in circulation many months, if not many years, after viral clearance.

MATERIALS AND METHODS

Subjects and cell lines. The generation of CD8 T-cell lines from the two women who were able to clear their HPV infections has been described previously (22). As a part of a longitudinal cohort study, they were monitored using cervical HPV DNA testing by PCR (27), cytology, and colposcopy every 4 months. Briefly, the CD8 T-cell lines were established by stimulating PBMCs using autologous dendritic cells (DCs) infected with recombinant vaccinia viruses expressing the HPV16 E6 protein. Using ELISPOT assays, the presence of potential CD8 T-cell epitopes in the E6 31–55 and E6 61–85 (dominant) regions was demonstrated for subject A (22) and that in the E6 31–55, E6 76–100, and E6 121–145 (dominant) regions was demonstrated for subject B (22). Each region was tested using a pool of three overlapping 15-mer peptides. The blood samples used to establish these CD8 T-cell lines were drawn 53 or 78 months after HPV16 clearance in subject A or B, respectively. The study protocol was approved by the University of California at San Francisco Committee on Human Research as well by the Institutional Review Board of the University of Arkansas for Medical Sciences.

Synthetic HPV peptides. A series of 15-mer peptides overlapping each other by 10 amino acids and a series of 9-mer peptides overlapping each other by 8 amino acids covering the HPV16 E6 protein have been described (20). To define the minimal and optimal amino acid sequences of the CD8 T-cell epitope, 8-mer, 10-mer, 11-mer, and homologous peptides (see Table 1) were synthesized as needed (CPC Scientific, Inc., San Jose, CA).

Isolation of antigen-specific T-cell clones after magnetic selection of IFN- γ -secreting cells. The CD8 T-cell lines from subjects A and B were stimulated for two additional 7-day cycles to increase the frequency of targeted antigen-specific T cells as described previously (20). Then, the HPV16 E6 antigen-specific CD8 T-cell lines ($\sim 2 \times 10^7$ cells from each line) were stimulated with 10 μ M of each peptide contained in positive peptide pools (three 15-mer peptides contained in each peptide pool) for 3 to 6 h. Following this incubation, the IFN- γ -secreting cells were selected using a commercially available kit according to the manufacturer's instructions (IFN- γ secretion assay; Miltenyi Biotec). HPV16 E6 peptide-specific T-cell clones were isolated using a limiting dilution method as described previously (20, 21).

ELISPOT assays for screening T-cell clones. A previously described ELISPOT assay method (21) was used to screen possible epitope-specific T-cell clones. In short, the plate (MultiScreen; Millipore, Bedford, MA) was coated with 5 μ g/ml anti-IFN- γ monoclonal antibody 1-DIK (Mabtech, Stockholm, Sweden) overnight at 4°C. After the plate was washed and blocked, 1×10^5 autologous Epstein-Barr virus-transformed B-lymphoblastoid cell line (LCL) cells were plated along with media containing T-cell clone cells (cell number not determined) to one well at the same position in triplicate (to test two peptide pools and a medium-only control for subject A) or quadruplicate (to test three peptide pools and a medium-only control for subject B) plates. The final concentration was 10 μ M for each peptide. After 20 h of incubation, the wells were washed and incubated with biotin-conjugated anti-IFN- γ monoclonal antibody (1 μ g/ml; Mabtech) for 2 h at 37°C. Then, the wells were washed and incubated with avidin-bound biotinylated horseradish peroxidase H (Vectastain Elite kit; Vector Laboratories, Inc., Burlingame, CA) for 1 h at 37°C. After additional washing of the wells, spots were developed using stable diaminobenzene (Research Genetics, Huntsville, AL) at room temperature. Spot-forming units were counted using an automated ELISPOT analyzer (Cell Technology, Inc., Jessup, MD). The wells that showed spots in an ELISPOT plate with one peptide pool, but not in other plates, were considered to potentially contain T-cell clones with the specificity of interest.

ELISPOT assays to characterize the CD8 T-cell epitopes of the HPV16 E6 protein. To confirm the specificity of the potential epitope-specific T-cell clones identified in screening, ELISPOT assays were repeated using 15-mer peptides contained in the positive peptide pools individually in duplicate or triplicate. One thousand T-cell clone cells were coincubated with 1×10^5 autologous LCL cells along with 20 U/ml of recombinant human interleukin-2 (rhIL-2) in the presence of 15-mer peptide at a concentration of 10 μ M. The ELISPOT assays were otherwise performed as described above. To determine the minimal and optimal amino acid sequences of the CD8 T-cell epitopes, additional ELISPOT assays were performed using peptides of various lengths. Serial dilutions of these peptides (10^{-5} M to 10^{-10} M) were also performed whenever necessary as

described previously (21). A peptide of particular length was considered optimal if noticeably larger numbers of spot-forming units were observed at multiple concentrations.

To determine whether the HPV16 E6 epitopes were being endogenously processed, autologous LCL cells infected with recombinant vaccinia virus expressing the E6 protein (E6-vac) at a multiplicity of infection of 5 were used as antigen-presenting cells (APCs) in ELISPOT assays. The wild-type virus, Western Reserve (WR), and/or recombinant vaccinia virus expressing HPV16 E7 (E7-vac) served as negative controls. The ELISPOT assays were otherwise performed as described above.

HLA typing. HLA typing was performed at the University of California at San Francisco Immunogenetics Laboratory or at the University of Arkansas for Medical Sciences HLA Laboratory by a serological method or a PCR sequence-specific amplification method.

Identification of the restricting HLA class I molecules. To identify putative restricting HLA class I molecules, ELISPOT assays were performed, as described above, using allogeneic LCLs sharing one or a few class I molecules with the subjects. The results were confirmed using chromium release assays in which at least two LCLs expressing the putative HLA class I molecule were tested. The LCL cells were labeled with 100 μ Ci of sodium chromate ($\text{Na}_2^{51}\text{CrO}_4$) and were incubated with 10 μ M of positive peptide (HPV16 E6 75–83 for subject A and E6 131–145 for subject B). After the cells were plated in a 96-well plate at 3×10^3 /well, effector cells were added at multiple effector/target cell (E:T) ratios. After a 5-h incubation, the supernatants were harvested and radioactivity was counted with a gamma counter (Packard Instruments, Meriden, CT). The percentage of specific lysis was calculated as previously described (23).

Characterization of the surface phenotypes of the T-cell clones. For characterization of the surface phenotypes of the T-cell clones, 5×10^5 T-cell clone cells were stained with CD4-phycoerythrin (PE)/CD8-fluorescein isothiocyanate (FITC) cocktail, CD3-FITC/CD16-PE cocktail, and corresponding isotype controls (Caltag, Burlingame, CA) for 30 min at 4°C. Then, the cells were washed and fixed. Events were acquired and analyzed using the Coulter EPICS XL-MLC flow cytometer (Beckman Coulter, Fullerton, CA).

Examination of the recognition of homologous CD8 T-cell epitopes from other high-risk HPV types using ELISPOT assays. Homologous epitopes, defined as peptides containing the same anchor residues (amino acid 2 and the last amino acid residue of the CD8 T-cell epitope) located within the 20-amino-acid region from the original HPV16 epitope, were identified by examining the protein sequences of the E6 proteins of HPV31, -33, -35, -39, -45, -51, -52, -56, -58, -59, -68, and -73 (HPV Sequence Database, Los Alamos National Laboratory, University of California). Recognition of the homologous epitopes by the HPV16 E6 75-83- or E6 133-142-specific T-cell clone cells (1,000 cells per well) was tested using autologous LCL cells and LCL cells (1×10^5 LCL cells per well) from five additional individuals expressing the HLA-B62 (for subject A) or HLA-A68 (for subject B) molecule. The HPV16 E6 75-83 and E6 133-142 peptides were used as positive controls. Peptides were added at a concentration of 10 μ M along with 20 U/ml of rhIL-2, and the wells were set up in duplicate.

Detection of epitope-specific memory T cells by the ELISPOT assay or tetramer analysis. During the process of stimulating and isolating epitope-specific T-cell clones, the CD8 T-cells from subjects A and B were incubated with autologous mature DCs infected with E6-vac since DCs are the most potent APCs. One drawback of this approach is that one cannot be certain whether the antigen-specific T cells are primed during the course of their natural HPV16 infection in vivo or during the course of the in vitro stimulation. If the former scenario is true, then the epitope-specific memory T cells may be detectable in PBMCs from the same subjects. In order to assess this possibility, a tetramer designed to bind T cells specific for the HPV16 E6 133–142 epitope restricted by the HLA-A6801 molecule was obtained from the NIH tetramer facility for subject B. A tetramer could not be made for the HPV16 E6 75–83 epitope restricted by the HLA-B62 molecule for subject A, since this HLA type was not available for production, and an ELISPOT assay was performed for the detection of memory T cells.

PBMCs from subject A (collected at 29 and 37 months) and from subject B (collected at 4, 22, 27, 48, and 74 months) after HPV16 clearance were available (Fig. 1). These PBMC samples were analyzed immediately after thawing (for subject B) and after in vitro stimulation in a manner known to enhance the number of memory T cells (10, 11, 15) (for subjects A and B). In short, cryopreserved PBMCs were thawed and incubated at 2×10^6 /ml in medium (RPMI 1640 plus 10% human serum, 100 U/ml penicillin, and 100 μ g/ml streptomycin; referred to as RP-10H) overnight. Then, the cells were counted and were incubated at 1×10^6 PBMCs/ml/well in the presence of the respective HPV16 E6 peptide at a concentration of 10 μ M (day 0). After a 3-day incubation, rhIL-2 was added to each well to a concentration of 1,800 U/ml. One more milliliter of




FIG. 1. Natural history of HPV16 infection. (a) For subject A, the HPV infection was detected at one visit. The T-cell clones isolated to define the dominant CD8 T-cell epitope were derived from a blood sample drawn 53 months after HPV clearance. PBMCs collected at 29 and 37 months after the clearance of HPV16 infection were available. Pos, positive; Neg, negative. (b) Subject B had an HPV16 infection which persisted for over 4 years. The T-cell clones isolated to define the dominant CD8 T-cell epitope were derived from a blood sample drawn 78 months after HPV16 clearance. PBMCs for tetramer staining were isolated from blood samples collected at 4, 22, 27, 48, and 74 months after the clearance of HPV16 infection.

RP-10H with rhIL-2 (1,800 U/ml) was added to each well on day 7. Three days later (day 10), the cells were washed and cultured in RP-10H at 10^6 /ml without peptide and rhIL-2 overnight. On the following day, 2.5×10^4 PBMCs per well were used for an ELISPOT assay as described above for subject A. The percentage of epitope-specific T cells was calculated by subtracting the averaged spot-forming units of no-peptide control wells from those of peptide-containing wells and by dividing the difference by the total number of cells.

For subject B, various tetramer concentrations, incubation times, and incubation temperatures were examined. The concentration of 1:1,600 and incubation time of 30 min at room temperature were chosen. In addition, the PBMC samples were stained with CD4-FITC, CD14-FITC, CD19-FITC, CD8-PerCP (peridinin-chlorophyll-protein complex), and CD45RO-allophycocyanin (BD Biosciences, San Jose, CA). PBMCs from a healthy donor known to be HLA-A6801 negative were used as a negative control. T-cell clone cells (#83-B designates clone 83 from subject B) were mixed with PBMCs from this negative control donor (10% T-cell clone cells) and were used as the positive control.

On day 11, the samples were stained and analyzed using Becton Dickinson FACSCalibur (BD Bioscience). A lymphocyte gate was drawn in forward and side scatter, and this population was further gated for CD4/14/19-FITC negativity to eliminate nonspecific staining of CD4 T cells, monocytes, and B cells. One-hundred thousand events were acquired per sample. The percentages of tetramer-positive CD8⁺ T cells were calculated by dividing the number of cells in the circle in the right upper corner (region determined using the positive control sample) by the sum of cell numbers in the upper right and lower right quadrants (CD8 T cells). Whether these tetramer-positive CD8⁺ T cells were CD45RO⁺ T cells was also determined.

RESULTS

Natural history of HPV16 infection in subjects A and B.

Subjects A and B were participants of a longitudinal study of HPV infection (18), and subject A's HPV16 infection was detected at a single visit, and the blood used to establish the CD8 T-cell line and to isolate T-cell clones was drawn 53 months later. On the other hand, subject B had an HPV16 infection which lasted for over 4 years, and the blood sample was taken 78 months after clearance (Fig. 1).

HPV16 E6 71–85 restricted by the HLA-B62 molecule; the dominant epitope characterized from subject A. Approximately 1.8×10^4 (0.09%) IFN- γ -secreting cells were isolated from the CD8 T-cell line from subject A. Limiting dilution analysis was performed, and a total of 344 T-cell clones were expanded. A random selection of 94 T-cell clones were used for screening ELISPOT, and eight T-cell clones (# 8-A, #15-A, #40-A, #74-A, #76-A, #78-A, #93-A, and #94-A)

were positive for the peptide pool covering the HPV16 E6 61–85 region (dominant region) (22), but none of the T-cell clones was positive for the peptide pool covering the E6 31–55 region (subdominant region) (22; data not shown). When the ELISPOT assay was repeated using individual 15-mer peptides, seven (#8-A, #15-A, #40-A, #74-A, #76-A, #78-A, and #94-A) of the eight screen-positive T-cell clones were positive with the E6 71–85 peptide but not with the E6 61–75 and E6 66–80 peptides (Fig. 2a). To examine the nature of the antigen processing, E6-vac- or WR-infected autologous LCL cells were used as APCs. The same seven T-cell clones were positive when tested with E6-vac-infected autologous LCL cells, suggesting that this E6 epitope is endogenously processed (Fig. 2b).

To determine the minimal and optimal epitope amino acid sequence, a series of ELISPOT assays were performed (Fig. 2c to e). Of the 9-mer peptides within the E6 71–85 region, the E6 75–83 9-mer peptide showed the most spot-forming units per well for all clones tested, followed by the E6 74–82 peptide (Fig. 2c). When the two 10-mer peptides surrounding the E6 75–83 peptide and two 8-mer peptides within were compared, the E6 75–83 9-mer peptide and the E6 74–83 10-mer peptide demonstrated comparable numbers of spot-forming units for all T-cell clones tested (Fig. 2d). The E6 74–83 9-mer peptide and the E6 75–84 10-mer peptide demonstrated considerably fewer spot-forming units. A serial dilution of the E6 75–83 9-mer peptide and the E6 74–83 10-mer peptide tested with clones #76-A and #78-A showed that they were equivalent over a wide range of peptide concentrations (Fig. 2e). Therefore, we designated E6 75–83 (9 amino acids) to be the minimal and optimal sequence for this epitope.

To identify the HLA restriction element of this novel epitope, a panel of allogeneic LCL cells matched to one or more HLA class I molecule(s) of subject A were used in an ELISPOT assay. An allogeneic LCL matched with the HLA-B62 molecule showed a positive response along with the autologous LCL cells for all four clones tested (#40-A, #76-A, #78-A, and #94-A) (data not shown). To confirm this result, a chromium release assay was performed with the T-cell clones (clones #76-A and #78-A) as effectors and LCL cells pulsed




FIG. 2. Characterization of subject A's dominant CD8 T-cell epitope in terms of its minimal and optimal amino acid sequence and the HLA-restricting molecule. The bars represent standard errors of the means. (a) ELISPOT assay performed to determine the specificity of the screen-positive T-cell clones to one of the three 15-mer peptides contained in the HPV16 E6 61–85 region. PHA, phytohemagglutinin. (b) ELISPOT assay performed to determine whether the epitope is endogenously presented using APCs infected with E6-vac or WR. (c) ELISPOT assay performed to identify the epitope within the 15-amino-acid region using 9-mer peptides overlapping by 8 amino acids. (d) ELISPOT assay performed to determine the optimal epitope sequence using two 10-mer peptides surrounding the E6 75–83 9-mer and two 8-mer peptides within it. The E6 74–82 9-mer peptide was also tested. (e) ELISPOT assay performed using serially diluted E6 75–83 9-mer peptide and E6 74–83 10-mer peptide (10^{-5} M to 10^{-10} M) to determine the optimal peptide of minimum length. A representative (clone #78-A) of two T-cell clones tested is shown. (f) Chromium release assay performed to determine the restricting molecule for the HPV16 E6 75–83 epitope using peptide-pulsed autologous LCL cells or a panel of partially HLA-matched LCL cells as APCs. A representative (#78-A) of the two clones tested is shown. θ , autologous LCL cells. *, HLA type determined using one of the molecular methods.

with the E6 75–83 peptide as targets. Three of the LCL cells were from allogeneic LCLs expressing the HLA-B62 molecule, and all of them were lysed by the T-cell clones (Fig. 2f), confirming that the restriction element of the HPV16 E6 75–83 epitope is the HLA-B62 molecule.

HPV16 E6 133–142 restricted by the HLA-A6801 molecule; the dominant epitope characterized from subject B. From this subject, 6.5×10^4 (0.31%) IFN- γ -secreting cells were selected, and 504 of 1,048 T-cell clones that grew were expanded. A ran-

dom selection of 94 T-cell clones was screened using ELISPOT. Sixty-four of 94 T-cell clones tested were positive for the E6 121–146 region (dominant region) (22), and none of them was positive for the E6 31–55 and E6 76–100 regions (subdominant regions) (22; data not shown). Eight (#1-B, #4-B, 13-B8, #21-B, #74-B, #80-B, #83-B, and #87-B) of 64 screen-positive T-cell clones that grew well were retested with three individual 15-mer peptides in the region as well as with autologous LCL cells infected with E6-vac, E7-vac, or WR. All




FIG. 3. Characterization of subject B's dominant CD8 T-cell epitope in terms of its minimal and optimal amino acid sequence and the HLA-restricting molecule. The bars represent standard errors of the means. (a) ELISPOT assay performed to determine the specificity of the screen-positive T-cell clones to one of the three 15-mer peptides contained in the HPV 16 E6 121–145 region. PHA, phytohemagglutinin. (b) ELISPOT assay performed to determine whether the epitope is endogenously presented using APCs infected with E6-vac or WR. (c) ELISPOT assay performed to identify the epitope within the 15-amino-acid region using 10-mer peptides overlapping by 9 amino acids. (d) ELISPOT assay performed to determine the optimal epitope sequence using two 11-mer peptides surrounding the E6 133–142 10-mer peptide and two 9-mer peptides within it. (e) ELISPOT assay performed using the serially diluted E6 133–142 10-mer peptide, E6 132–142 11-mer peptide, and E6 133–142 11-mer peptide (10^{-5} M to 10^{-10} M) to determine the optimal peptide of minimal length. (f) Chromium release assay performed to determine the restricting molecule for the HPV16 E6 133–142 epitope using peptide-pulsed autologous LCL cells or a panel of partially HLA-matched LCL cells as APCs. A representative (#87-B) of the two clones tested is shown. θ , autologous LCL cells. *, HLA type determined using one of the molecular methods.

eight T-cell clones were positive with the E6 131–145 peptide and E6-vac (Fig. 3a and b). As with the previous subject, a series of ELISPOT assays were performed to determine the optimal peptide of minimal length (Fig. 3c to e). Since none of the 9-mer peptides included in the E6 131–145 region was positive (data not shown), the ELISPOT assay was repeated with a series of 10-mer peptides, and only E6 133–142 was positive among the 10-mers (Fig. 3c). When two 11-mers surrounding this E6 133–142 10-mer and two 9-mers within it were tested, the 10-mer and the two 11-mers appeared equivalent (Fig. 3d). This was true even over a wide range of concentrations (Fig. 3e); therefore, the E6 133–142 10-mer peptide was designated to be the optimal peptide of minimal length.

A panel of allogeneic LCL cells matched to one or more HLA class I molecule(s) of subject B were used in an ELISPOT assay, and allogeneic LCL cells matched with the HLA-A68 molecule showed a positive response along with the autologous LCL cells for all eight clones tested (#1-B, #4-B, #13-B, #21-B, #74-B, #80-B, #83-B, and #87-B) (data not shown). To confirm this result, a chromium release assay was performed with the T-cell clones (clones #83-B and #87-B) as effectors and LCL cells pulsed with the E6 131–145 peptide as targets. Of the three HLA-A68-positive LCL cells, one was known to be HLA-A6801 while another one was known to be HLA-A6802 (Fig. 3f). The results were positive for autologous LCL cells and for LCL cells expressing A68 or A6801. Other allogeneic

TABLE 1. Amino acid sequences of high-risk HPV type homologous epitopes homologous for HPV16 E6 75–83 and E6 133–142 CD8 T-cell epitopes

CD8 T-cell epitope homology and HPV type	Amino acid sequence ^a	Position	Sequence length (no. of amino acids)
Homologous for HPV 16 E6 75–83 epitopes			
HPV16	KFYSKISEY	75–83	9
HPV33	RFLSKISEY	68–76	9
HPV51	LFYSKIREY	68–76	9
HPV52	RFLSKISEY	68–76	9
HPV56	LFYSKVRKY	71–79	9
HPV73	KFYSKIREY	69–77	9
Homologous for HPV16 E6 133–142 epitopes			
HPV16	HNIRGRWTGR	133–142	10
HPV31	HNIGGRWTGR	126–135	10
HPV33	HNISGRWAGR	126–135	10
HPV51	ANCWQTRQR	137–146	10
HPV52	HNIMGRWTGR	126–135	10
HPV58	HNISGRWTGR	126–135	10

^a Boldface amino acid residues are different from those in HPV16 E6 75–83 or E6 133–142 CD8 T-cell epitope.

LCL cells were negative, including the allogeneic LCL cells expressing the HLA-6802 molecule. Taken together, these data suggest that the restriction element of the HPV16 E6 133–142 epitope is the HLA-A6801 molecule.

No recognition of homologous epitopes from other high-risk HPV by the HPV16 E6 75-83-specific and E6 133-142-specific T-cell clone cells. To determine the potential recognition by epitope-specific T-cell clone cells, the presence of homologous epitopes from other high-risk HPV types were examined. Five homologous epitopes for the HPV16 E6 75–83 CD8 T-cell epitope (HPV33, -51, -52, -56, and -73) and five for the HPV16 133–142 epitope (HPV31, -33, -51, -52, and -58) were identified (Table 1). Their recognition by the HPV16 E6 75-83-specific T-cell clones (#76-A and #78-A) was examined by peptide-pulsed autologous LCL cells and five allogeneic LCL cells expressing the HLA-B62 molecule. All of the LCL cells were recognized, as measured by the number of spot-forming units in an ELISPOT assay when they were pulsed with the HPV16 E6 75-83 peptide but not with any other homologous epitopes (data not shown). The results were the same for the HPV16 E6 133-142 epitope in that the LCL cells were recognized only when pulsed with the HPV16 E6 133-142 peptide but not with the others (#83-B and #87-B) (data not shown). These data imply that the recognition of these HPV16 E6 epitopes is type specific.

HPV E6 75-83- and E6 133-142-specific T-cell clone cells show the surface phenotype of CD3⁺ CD4⁻ CD8⁺ CD16⁻. The surface phenotypes of the E6 75-83- and E6 133-142-specific T-cell clones have been demonstrated to be CD3⁺ CD4⁻ CD8⁺ CD16⁻ (#40-A, #76-A, #78-A, #94-A, #1-B, #4-B, #13-B, #21-B, #74-B, #80-B, #83-B, and #87-B) (data not shown).

The presence of epitope-specific memory T cells in subjects A and B. Peptide-stimulated PBMC samples from subject A

collected 29 and 37 months after HPV16 clearance (Fig. 1) demonstrated the frequencies of the epitope-specific T cells to be 0.039% and 0.092%, respectively (Fig. 4).

PBMC samples from subject B collected at 4, 22, 27, 48, and 74 months (Fig. 1) after the clearance of HPV16 infection were analyzed with and without in vitro stimulation with the HPV16 E6 133–142 peptide (Fig. 5 and Table 2). Three (one tetramer-positive CD8⁺ T-cell in 2,146 CD8 T cells at 4 months, one in 3,116 at 22 months, and one in 2,311 at 74 months) of the five PBMC samples (without in vitro stimulation) demonstrated staining for the tetramer-positive CD8⁺ T cells above that of the negative control (1 in 5,010). Upon stimulation with the HPV16 E6 133–142 peptide, the frequencies of these tetramer-positive CD8⁺ T cells increased noticeably in the PBMC samples taken at 4 months and 22 months (1 in 512 and 1 in 510, respectively). Ninety-eight percent and 100% of these the tetramer-positive CD8⁺ T cells were CD45RO⁺ in these samples, respectively (data not shown). Taken together, the HPV16 E6 133-142-specific memory T-cells were detectable at 4 months and 22 months after HPV16 clearance.

DISCUSSION

In the present study, we characterized HPV16-specific CD8 T-cell epitopes from two women in whom the presence of potential CD8 T-cell epitopes was demonstrated after HPV16 clearance (22). For one subject (subject A), the duration of the HPV16 infection was brief and it was detectable only at a single visit, while that of the other subject (subject B) lasted for over 4 years (Fig. 1). Although both subjects demonstrated the presence of potential HPV16 E6 CD8 T-cell epitopes within more than one region, only the dominant T-cell epitopes were characterized since we were unsuccessful in isolating T-cell clones from the subdominant regions. These dominant epitopes were characterized to be the HPV16 E6 75–83 epitope restricted by the HLA-B62 molecule and the HPV16 E6 133–142 epitope restricted by the HLA-A6801 molecule. The antigen frequencies of the HLA-B62 molecule are 0.9%, 6.2%, and 18.1% in African, Caucasian, and Asian populations, respectively, and those of the HLA-A68 molecule are 19.4%, 8.0%, and 2.6% in the respective populations (17). Previously, we had characterized HPV16-specific CD8 T-cell epitopes from the same cohort of women (20, 22) and described the phenom-




FIG. 4. ELISPOT assay performed to determine the frequencies of the epitope-specific T cells after clearance of HPV16 infection but prior to isolation of the T-cell clones for subject A. The bars represent standard errors of the means.




FIG. 5. Tetramer staining performed to determine frequencies of epitope-specific T cells after clearance of HPV16 infection but prior to isolation of T-cell clones for subject B. Cells were selected using a lymphocyte gate in forward and side scatter and for being CD4/14/19 negative. (a) Tetramer staining of PBMCs from a healthy donor known to be HLA-A6801 negative mixed with HPV16 E7 133–142 T-cell clone cells (#83-B) at a 10:1 ratio (positive control). (b) Tetramer staining of PBMCs from a healthy donor known to be HLA-A6801 negative (negative control). (c) Tetramer staining of peptide-stimulated PBMC sample collected 4 months after the clearance of HPV infection. (d) Tetramer staining of peptide-stimulated PBMC sample collected 22 months after the clearance.

enon of HLA class I binding promiscuity among the newly described epitopes (21). Similarly, the HPV16 E6 75–83 epitope has an overlap with another CD8 T-cell epitope, HPV16 E6 80–88, previously described by Bourgault Villada and colleagues (3). To our knowledge, there are no other CD8 T-cell epitopes that overlap with the HPV16 E6 132–142 epitope. However, Gallagher and Man have recently described a CD4 T-cell epitope, HPV16 E6 127–141, restricted by the HLA-DRB1*01 (DR1) molecule (8). Therefore, it would be possible to develop T-cell epitope-based HPV vaccines which include these epitopes by covering small parts of the HPV16 E6 protein: i.e., the HPV16 E6 75–88 and E6 127–142 regions.

TABLE 2. Frequencies of HPV16 E6 133–142 tetramer-positive CD8⁺ T cells among CD8⁺ T cells

No. of mo after HPV clearance or control type	Result for PBMCs:			
	Untreated		Peptide stimulated ^a	
	% Tetramer ⁺ CD8 ⁺	Reciprocal frequency	% Tetramer ⁺ CD8 ⁺	Reciprocal frequency
4	0.047	2,146	0.20	512
22	0.032	3,116	0.20	510
27	0.020	5,002	0.0086	11,572
48	0.019	5,214	0.0052	19,350
74	0.043	2,311	0.0098	10,209
Control				
Positive	9.0	11	12.6	8
Negative	0.020	5,010	0.0084	11,939

^a PBMCs were stimulated with HPV16 E6 133–142 for 10 days before analysis.

The utility of the newly described HPV16 E6 CD8 T-cell epitopes as the source of antigens for peptide-based vaccine or immunotherapy would be broader if the epitope-specific T cells also recognized homologous epitopes from other high-risk HPV types. To evaluate whether these two newly described CD8 T-cell epitopes may be similar enough to potential epitopes from other high-risk HPV types, recognition of homologous epitopes (Tables 1) by the HPV16 E6 75-83- or E6 132-142-specific T-cell clones was examined. For these epitopes, no recognition was demonstrated, suggesting that these two T-cell epitopes are type specific. These results are in contrast to the HPV16 E6 52–61 epitope, which was evaluated by the same method, in which the recognition of homologous epitopes from HPV35, -39, -45, -51, and -73 by HPV16 E6 52-61-specific T-cell clone cells was shown (16).

Memory T cells play an important role in maintaining long-term immunity to previously encountered pathogens or tumor antigens. They may proliferate, rapidly acquire effector functions to kill virus-infected cells or tumor cells, or secrete cytokines that inhibit replication of the pathogen after restimulation with reexposure to antigen (14). For subject A, the epitope-specific memory T cells were detected in circulation 37 months after viral clearance using an ELISPOT assay (Fig. 4). For subject B, tetramer-positive CD8⁺ T cells were present *ex vivo* in three of the five PBMC samples (4, 22, and 74 months). Upon peptide stimulation *in vitro* for 10 days, the frequencies of tetramer-positive CD8⁺ T cells expanded in two of the three PBMC samples (4 and 22 months; Table 2), and almost all of the epitope-specific T cells were CD45RO⁺. There is no good

explanation as to why PBMCs collected at 74 months were not confirmed to be positive after peptide stimulation, while those collected at 4 and 22 months were. One possibility is that there is not a sufficient difference in the frequencies of the tetramer-positive PBMCs (0.043% at 74 months) and the negative control (0.020%), such that the samples that appear positive (4, 22, and 74 months) are actually in a gray zone between positive and negative. This is a reflection of a low number of circulating HPV-specific CD8 T cells, especially long after viral clearance. Since this method of *in vitro* stimulation is used to increase the number of memory T cells (10, 11, 15), the data seem to indicate that the HPV-specific memory T cells can certainly be detected up to 37 months, or 3 years, after the HPV DNA has become undetectable. Furthermore, the fact that epitope-specific T memory cells were detected prior to isolation of T-cell clones suggests that they were generated *in vivo* at the time of HPV16 infection rather than *in vitro*. It may be that the HPV16 E6 132-142-specific T cells were isolated 78 months after HPV16 clearance because mature DCs, which are known to be the most potent APCs, were able to amplify the number of epitope-specific memory T cells to a level at which they can be isolated. Godkin et al. demonstrated a memory T-cell response to hepatitis virus epitopes many years after viral clearance (10), and Demkowicz et al. showed persistence of T cells specific for vaccinia virus for up to 50 years in the absence of reexposure (5). The challenge in the future is to determine whether these memory cells confer long-term protection to HPV reinfection and whether these memory T cells are capable of quickly differentiating into a potent effector response during HPV reinfection.

Tetramers specific for HPV16 E6 or E7 CD8 T-cell epitopes have been used by other investigators to examine the frequencies of HPV-specific T lymphocytes in patients with cervical cancer (30, 31), with cervical intraepithelial neoplasia grade 3 (CIN 3) (30), and with squamous cell carcinoma of the oropharynx (SCCO) (13). In T-cell lines established from five HLA-A*0201-positive patients with HPV16-positive cervical cancer by stimulation with the HPV16 E7 11–20 peptide, 1 to 12% of cells were tetramer positive (30). However, *ex vivo* analysis of PBMCs from patients with CIN 3 (1 of 1,260 to 1 of 19,073) and normal controls (1 of 1,855 to 1 of 42,004) revealed low frequencies of tetramer-positive CD8⁺ cells in both. Similar findings were reported by Hoffmann and colleagues in that the frequencies of tetramer-positive CD8⁺ cells for three epitopes (HPV16 E7 11–20, E7 82–90, and E7 86–93) studies were not significantly different between patients with SCCO and healthy controls (13). However, a significant difference was found in frequencies of the HPV16 E7 11-20-specific T cells between patients with SCCO whose tumors were positive for the expression of HPV16 E7 and p16 and who were negative for these two proteins. Zehbe et al. examined the presence of HPV16 E6 11-29-specific T cells in PBMCs, tumor-infiltrating lymphocytes, and in lymph nodes of five HLA-A*0201-positive patients with HPV16-positive cervical cancer (31). Interestingly, the percentage of tetramer-positive CD8⁺ cells was highest in the tumor-infiltrating lymphocytes but they had a naïve phenotype (CD45RA⁺ CCR7⁺). To our knowledge, our group is the first to examine the presence of HPV-specific T lymphocytes using a tetramer in a woman whose HPV16 infection and its clearance were documented.

Vaccination with HPV16 E7 CTL epitopes was shown to elicit efficient antitumor CTL responses and to prevent the outgrowth of a lethal dose of HPV16-induced tumor cells in mice, indicating the feasibility of peptide immunization (7, 25). However, the use of peptides as HPV therapeutic vaccines in human clinical trials has shown only limited success. Rensing et al. (26) used two CTL epitopes, HPV16 E7 11–20 and E7 86–93, and a universal T-helper epitope to perform a peptide-based phase I/II vaccination trial to induce antitumor immune responses in patients with recurrent or residual cervical carcinoma. Though no HPV16-specific CTL response was detectable, strong T-helper peptide-specific proliferations were detected in 4 of 12 patients, showing T-helper epitopes may also play a role in the generation of protective antitumor immunity. In another trial, 18 women with high-grade cervical or vulvar intraepithelial neoplasia II/III who were positive for HPV16 received vaccination with synthetic E7 12–20 peptide representing the CTL epitope (19). The E7 86–93 peptide linked to a helper T-cell epitope peptide was administered to part of the women at the same time. It was demonstrated that six patients had partial colposcopically measured regression of their CIN lesions, and three patients completely cleared their dysplasia after vaccine. Similarly, an epitope-based HPV therapeutic vaccine delivered in a form of a DNA vaccine has shown partial success. ZYC101a contains a plasmid encoding portions derived from the HPV16 and -18 E6 and E7 proteins. In a phase II clinical trial that enrolled 127 evaluable subjects with CIN 2/3 (9), a higher percentage of subjects receiving ZYC101a demonstrated regression compared to placebo (43% versus 27%). However, the difference was not statistically significant ($P = 0.12$). In a subgroup of women ≤ 25 years old ($n = 43$), the regression rate was significantly higher in the vaccine group than in the placebo group (70% versus 23%; $P = 0.007$). In short, we described two novel HPV16 E6 CD8 T-cell epitopes from women who were able to clear their HPV16 infection and demonstrated the presence of HPV-specific memory T cells years after its clearance. The challenges for the future, in order to translate these findings to the development of effective vaccine, are to evaluate more efficient methods of presenting the T-cell epitopes so the clinical response rates would be more robust.

ACKNOWLEDGMENTS

We thank the subjects for participation in the study and Zhixiang Qu for technical expertise.

This work was supported by the National Institutes of Health (NCI CA051323) and the American Cancer Society Scholars Award (RSG-06-180-01-MBC).

REFERENCES

- Alexander, M., M. L. Salgaller, E. Celis, A. Sette, W. A. Barnes, S. A. Rosenberg, and M. A. Steller. 1996. Generation of tumor-specific cytolytic T lymphocytes from peripheral blood of cervical cancer patients by *in vitro* stimulation with a synthetic human papillomavirus type 16 E7 epitope. *Am. J. Obstet. Gynecol.* **175**:1586–1593.
- Bontkes, H. J., T. D. de Gruijl, J. M. Walboomers, A. J. van den Muysenberg, A. W. Gunther, R. J. Scheper, C. J. Meijer, and J. A. Kummer. 1997. Assessment of cytotoxic T-lymphocyte phenotype using the specific markers granzyme B and TIA-1 in cervical neoplastic lesions. *Br. J. Cancer* **76**:1353–1360.
- Bourgault Villada, I., N. Beneton, C. Bony, F. Connan, J. Monsonego, A. Bianchi, P. Saiag, J. P. Levy, J. G. Guillet, and J. Choppin. 2000. Identification in humans of HPV-16 E6 and E7 protein epitopes recognized by cytolytic T lymphocytes in association with HLA-B18 and determination of the HLA-B18-specific binding motif. *Eur. J. Immunol.* **30**:2281–2289.

4. Crook, T., J. A. Tidy, and K. H. Vousden. 1991. Degradation of p53 can be targeted by HPV E6 sequences distinct from those required for p53 binding and trans-activation. *Cell* **67**:547–556.
5. Demkowicz, W. E., Jr., R. A. Littaua, J. Wang, and F. A. Ennis. 1996. Human cytotoxic T-cell memory: long-lived responses to vaccinia virus. *J. Virol.* **70**:2627–2631.
6. Evans, E. M., S. Man, A. S. Evans, and L. K. Borysiewicz. 1997. Infiltration of cervical cancer tissue with human papillomavirus-specific cytotoxic T-lymphocytes. *Cancer Res.* **57**:2943–2950.
7. Feltkamp, M. C., H. L. Smits, M. P. Vierboom, R. P. Minnaar, B. M. de Jongh, J. W. Drijfhout, J. ter Schegget, C. J. Melief, and W. M. Kast. 1993. Vaccination with cytotoxic T lymphocyte epitope-containing peptide protects against a tumor induced by human papillomavirus type 16-transformed cells. *Eur. J. Immunol.* **23**:2242–2249.
8. Gallagher, K. M., and S. Man. 2007. Identification of HLA-DR1- and HLA-DR15-restricted human papillomavirus type 16 (HPV16) and HPV18 E6 epitopes recognized by CD4⁺ T cells from healthy young women. *J. Gen. Virol.* **88**:1470–1478.
9. Garcia, F., K. U. Petry, L. Munderspach, M. A. Gold, P. Braly, C. P. Crum, M. Magill, M. Silverman, R. G. Urban, M. L. Hedley, and K. J. Beach. 2004. ZYC101a for treatment of high-grade cervical intraepithelial neoplasia: a randomized controlled trial. *Obstet. Gynecol.* **103**:317–326.
10. Godkin, A. J., H. C. Thomas, and P. J. Openshaw. 2002. Evolution of epitope-specific memory CD4⁺ T cells after clearance of hepatitis C virus. *J. Immunol.* **169**:2210–2214.
11. Goonetilleke, N., S. Moore, L. Dally, N. Winstone, I. Cebere, A. Mahmoud, S. Pinheiro, G. Gillespie, D. Brown, V. Loach, J. Roberts, A. Guimaraes-Walker, P. Hayes, K. Loughran, C. Smith, J. De Bont, C. Verlinde, D. Vooijs, C. Schmidt, M. Boaz, J. Gilmour, P. Fast, L. Dorrell, T. Hanke, and A. J. McMichael. 2006. Induction of multifunctional human immunodeficiency virus type 1 (HIV-1)-specific T cells capable of proliferation in healthy subjects by using a prime-boost regimen of DNA- and modified vaccinia virus Ankara-vectored vaccines expressing HIV-1 Gag coupled to CD8⁺ T-cell epitopes. *J. Virol.* **80**:4717–4728.
12. Greenberg, P. D. 1991. Adoptive T cell therapy of tumors: mechanisms operative in the recognition and elimination of tumor cells. *Adv. Immunol.* **49**:281–355.
13. Hoffmann, T. K., C. Arsov, K. Schirlau, M. Bas, U. Friebe-Hoffmann, J. P. Klussmann, K. Scheckenbach, V. Balz, H. Bier, and T. L. Whiteside. 2006. T cells specific for HPV16 E7 epitopes in patients with squamous cell carcinoma of the oropharynx. *Int. J. Cancer* **118**:1984–1991.
14. Kaech, S. M., E. J. Wherry, and R. Ahmed. 2002. Effector and memory T-cell differentiation: implications for vaccine development. *Nat. Rev. Immunol.* **2**:251–262.
15. Keating, S. M., P. Bejon, T. Berthoud, J. M. Vuola, S. Todryk, D. P. Webster, S. J. Dunachie, V. S. Moorthy, S. J. McConkey, S. C. Gilbert, and A. V. Hill. 2005. Durable human memory T cells quantifiable by cultured enzyme-linked immunospot assays are induced by heterologous prime boost immunization and correlate with protection against malaria. *J. Immunol.* **175**:5675–5680.
16. Kim, K. H., R. Dishongh, A. D. Santin, M. J. Cannon, S. Bellone, and M. Nakagawa. 2006. Recognition of a cervical cancer derived tumor cell line by a human papillomavirus type 16 E6 52-61-specific CD8 T cell clone. *Cancer Immunol.* **6**:9.
17. Marsh, S. G. E., P. Parham, and L. D. Barber. 2000. *The HLA FactsBook*. Academic Press, San Diego, CA.
18. Moscicki, A. B., S. Shiboski, J. Broering, K. Powell, L. Clayton, N. Jay, T. M. Darragh, R. Brescia, S. Kanowitz, S. B. Miller, J. Stone, E. Hanson, and J. Palefsky. 1998. The natural history of human papillomavirus infection as measured by repeated DNA testing in adolescent and young women. *J. Pediatr.* **132**:277–284.
19. Munderspach, L., S. Wilczynski, L. Roman, L. Bade, J. Felix, L. A. Small, W. M. Kast, G. Fascio, V. Marty, and J. Weber. 2000. A phase I trial of a human papillomavirus (HPV) peptide vaccine for women with high-grade cervical and vulvar intraepithelial neoplasia who are HPV 16 positive. *Clin. Cancer Res.* **6**:3406–3416.
20. Nakagawa, M., K. H. Kim, and A.-B. Moscicki. 2004. Different methods of identifying new antigenic epitopes of human papillomavirus type 16 E6 and E7 proteins. *Clin. Diagn. Lab. Immunol.* **11**:889–896.
21. Nakagawa, M., K. H. Kim, T. M. Gillam, and A.-B. Moscicki. 2007. HLA class I binding promiscuity of the CD8 T-cell epitopes of human papillomavirus type 16 E6 protein. *J. Virol.* **81**:1412–1423.
22. Nakagawa, M., K. H. Kim, and A. B. Moscicki. 2005. Patterns of CD8 T-cell epitopes within the human papillomavirus type 16 (HPV 16) E6 protein among young women whose HPV 16 infection has become undetectable. *Clin. Diagn. Lab. Immunol.* **12**:1003–1005.
23. Nakagawa, M., D. P. Stites, S. Farhat, J. R. Sisler, B. Moss, F. Kong, A. B. Moscicki, and J. M. Palefsky. 1997. Cytotoxic T lymphocyte responses to E6 and E7 proteins of human papillomavirus type 16: relationship to cervical intraepithelial neoplasia. *J. Infect. Dis.* **175**:927–931.
24. Nakagawa, M., D. P. Stites, S. Patel, S. Farhat, M. Scott, N. K. Hills, J. M. Palefsky, and A. B. Moscicki. 2000. Persistence of human papillomavirus type 16 infection is associated with lack of cytotoxic T lymphocyte response to the E6 antigens. *J. Infect. Dis.* **182**:595–598.
25. Rensing, M. E., A. Sette, R. M. Brandt, J. Ruppert, P. A. Wentworth, M. Hartman, C. Oseroff, H. M. Grey, C. J. Melief, and W. M. Kast. 1995. Human CTL epitopes encoded by human papillomavirus type 16 E6 and E7 identified through in vivo and in vitro immunogenicity studies of HLA-A*0201-binding peptides. *J. Immunol.* **154**:5934–5943.
26. Rensing, M. E., W. J. van Driel, R. M. Brandt, G. G. Kenter, J. H. de Jong, T. Bauknecht, G. J. Fleuren, P. Hoogerhout, R. Offringa, A. Sette, E. Celis, H. Grey, B. J. Trimbos, W. M. Kast, and C. J. Melief. 2000. Detection of T helper responses, but not of human papillomavirus-specific cytotoxic T lymphocyte responses, after peptide vaccination of patients with cervical carcinoma. *J. Immunother.* **23**:255–266.
27. Ting, Y., and M. M. Manos. 1990. *Detection and typing of genital human papillomaviruses*. Academic Press, San Diego, CA.
28. Vousden, K. 1993. Interactions of human papillomavirus transforming proteins with the products of tumor suppressor genes. *FASEB J.* **7**:872–879.
29. Walboomers, J. M., M. V. Jacobs, M. M. Manos, F. X. Bosch, J. A. Kummer, K. V. Shah, P. J. Snijders, J. Peto, C. J. Meijer, and N. Munoz. 1999. Human papillomavirus is a necessary cause of invasive cervical cancer worldwide. *J. Pathol.* **189**:12–19.
30. Youde, S. J., P. R. Dunbar, E. M. Evans, A. N. Fiander, L. K. Borysiewicz, V. Cerundolo, and S. Man. 2000. Use of fluorogenic histocompatibility leukocyte antigen-A*0201/HPV 16 E7 peptide complexes to isolate rare human cytotoxic T-lymphocyte-recognizing endogenous human papillomavirus antigens. *Cancer Res.* **60**:365–371.
31. Zehbe, I., A. M. Kaufmann, M. Schmidt, H. Hohn, and M. J. Maeurer. 2007. Human papillomavirus 16 E6-specific CD45RA⁺ CCR7⁺ high avidity CD8⁺ T cells fail to control tumor growth despite interferon-gamma production in patients with cervical cancer. *J. Immunother.* **30**:523–532.
32. zur Hausen, H. 1999. immortalization of human cells and their malignant conversion by high risk human papillomavirus genotypes. *Semin. Cancer Biol.* **9**:405–411.
33. zur Hausen, H. 1996. Papillomavirus infections—a major cause of human cancers. *Biochim. Biophys. Acta* **1288**:F55–F78.