

Opsonophagocytic Assay To Evaluate Immunogenicity of Nontyphoidal *Salmonella* Vaccines

Girish Ramachandran,^{a,b} Mary Adetunke Boyd,^{a,c} Jennifer MacSwords,^{a,b} Ellen E. Higginson,^{a,b} Raphael Simon,^{a,b} James E. Galen,^{a,b} Marcela F. Pasetti,^{a,c} Myron M. Levine,^{a,b,c} Sharon M. Tennant^{a,b}

Center for Vaccine Development,^a Department of Medicine,^b and Department of Pediatrics,^c University of Maryland School of Medicine, Baltimore, Maryland, USA

Nontyphoidal *Salmonella* (NTS) invasive infections are an important cause of morbidity and mortality in sub-Saharan Africa. Several vaccines are in development to prevent these infections. We describe an NTS opsonophagocytic killing assay that uses HL-60 cells and baby rabbit complement to quantify functional antibodies elicited by candidate NTS vaccines.

Invasive nontyphoidal *Salmonella* (NTS) infections are increasingly being recognized as an important cause of morbidity and mortality in infants and HIV-infected adults and children in sub-Saharan Africa (1). Multiple vaccines are currently in development, including live-attenuated vaccines, conjugate vaccines, and generalized modules for membrane antigens against invasive *S. Typhimurium* and *S. Enteritidis* (2–7). As these vaccines enter clinical development, it is important to establish their immunogenicity through relevant, well-characterized immunological assays. There is a growing realization within the vaccinology community that in addition to determining antibody levels through binding assays, it is important to determine the functional, antimicrobial capacity of these antibodies.

Although many studies have evaluated anti-*Salmonella* serum bactericidal and opsonophagocytic antibody activity (2, 8–15), there is a need for reproducible assays that can be used routinely to characterize functional antibody responses to *Salmonella* in a standardized manner. We previously developed a complement-dependent serum bactericidal antibody (SBA) activity assay that quantifies serum antibody responses to typhoidal (*S. Typhi* and *S. Paratyphi A*) and nontyphoidal (*S. Typhimurium* and *S. Enteritidis*) *Salmonella* (16). Here, we describe an assay that measures the opsonophagocytic capacity of NTS antibodies based on the well-characterized pneumococcus opsonophagocytic activity (OPA), which employs HL-60 phagocytic cells and a standardized source of complement, i.e., baby rabbit complement (BRC) (17, 18).

The *Salmonella* OPA assay was evaluated using serum samples from mice that had been vaccinated with live-attenuated NTS vaccines (7). BALB/c mice were orally immunized with 10^9 CFU of *S. Typhimurium* CVD 1931 or *S. Enteritidis* CVD 1944 suspended in phosphate-buffered saline (PBS) or given PBS alone on days 0, 28, and 56, as previously described (7). *S. Typhimurium* CVD 1931 and *S. Enteritidis* CVD 1944 are vaccine strains derived from invasive disease-associated clinical isolates (*S. Typhimurium* D65 and *S. Enteritidis* R11, respectively), which harbor deletions in *guaBA* and *clpX*. Serum samples from vaccinated mice taken 83 days postimmunization were designated immune, and serum samples from mice that received PBS were designated nonimmune.

HL-60 cells were maintained in RPMI 1640 complete medium (RPMI 1640 [Corning CellGro, Manassas, VA] supplemented with 10% [vol/vol] fetal bovine serum [FBS; HyClone, Waltham, MA] and 1% GlutaMAX, 1% sodium pyruvate, and 1% penicillin-streptomycin [Life Technologies]) and differentiated for 6 days in

RPMI 1640 complete medium, 0.8% dimethylformamide (Fisher Scientific, Waltham, MA). Prior to use, cells were washed once in Hanks balanced salt solution (HBSS) (Life Technologies) and re-suspended to 1×10^6 cells/ml in opsonization buffer (OPB) (HBSS, 0.1% gelatin [Sigma-Aldrich, St. Louis, MO], 10% FBS). *Salmonella* strains (*S. Typhimurium* D65 and *S. Enteritidis* SO1 [19]) from overnight cultures were diluted 1 in 1,000 in Hy-Soy medium (0.5% sodium chloride, 1% Hy-Soy [Kerry, Clackmannshire, United Kingdom], and 0.5% Hy-Yest [Kerry]), grown at 37°C to an optical density at 600 nm (OD_{600}) of 0.3, and then diluted to 3×10^4 CFU/ml in OPB. Immune and nonimmune sera were heat-inactivated at 56°C for 20 min, and 2-fold serial dilutions in OPB (25 μ l/well final volume) were performed in a U-bottomed 96-well microplate (Sigma-Aldrich, St. Louis, MO). The bottom row was left without sera to act as a negative control. To each well, 3×10^2 CFU of bacterial suspension in 10 μ l OPB was added. Opsonization of bacteria was allowed to occur for 15 min at 37°C with 5% CO_2 . After incubation, 25 μ l of BRC (lot 31235; Pel-Freez Biologicals, Rogers, AZ) was added to the wells at a final concentration of 12.5% (vol/vol), and 40 μ l of differentiated HL-60 cells was added at a concentration of 4×10^4 cells/well. Plates were incubated at 37°C with shaking at 160 rpm for 45 min. Bacteria were enumerated by mixing the contents of each well by pipetting vigorously and performing viable counts in triplicate for each well. OPA titer was defined as the reciprocal of the highest serum dilution that produced >50% killing compared with the killing observed in the negative-control wells. Titers were determined from the mean bacterial counts in duplicate wells.

First, we determined the components required to demonstrate antibody-mediated OPA. Serum samples from mice immunized with CVD 1931 (*S. Typhimurium* D65 Δ *guaBA* Δ *clpX*) were

Received 29 February 2016 Accepted 23 March 2016

Accepted manuscript posted online 30 March 2016

Citation Ramachandran G, Boyd MA, MacSwords J, Higginson EE, Simon R, Galen JE, Pasetti MF, Levine MM, Tennant SM. 2016. Opsonophagocytic assay to evaluate immunogenicity of nontyphoidal *Salmonella* vaccines. *Clin Vaccine Immunol* 23:520–523. doi:10.1128/CVI.00106-16.

Editor: D. L. Burns, Food and Drug Administration

Address correspondence to Sharon M. Tennant, stennant@medicine.umaryland.edu.

G.R. and M.A.B. are co-first authors.

Copyright © 2016, American Society for Microbiology. All Rights Reserved.




FIG 1 Optimization of OPA assay. To determine all of the components required for optimal opsonophagocytosis, we optimized the OPA assay using pooled immune serum produced by live-attenuated *S. Typhimurium* CVD 1931 (*S. Typhimurium* D65 Δ *guaBA* Δ *clpX*) and measured the killing of *S. Typhimurium* D65. We determined the effect of combining HL-60 cells and baby rabbit complement (BRC) and immune sera (I) or nonimmune sera (NI) in various combinations. OPA titer was determined as the reciprocal of the highest serum dilution that produced >50% killing in comparison to bacteria incubated with BRC and HL-60 cells. The dashed line represents the lowest dilution tested. Results shown are the titers obtained from a single experiment. Titers obtained for the conditions containing immune serum were verified by repeating the experiment using a different pool of immune serum from the same mice. Similar OPA titers were obtained (I + HL-60 + BRC, 1:204,800; I + HL60, 1:100; and I + BRC, 1:100).

tested for OPA with HL-60 cells and BRC in different combinations against an invasive clinical isolate of *S. Typhimurium* D65 from Mali, as shown in Fig. 1. In the absence of HL-60 cells, there was no detectable bacterial killing. Thus, at a con-




FIG 2 OPA titers for individual mice immunized with *S. Typhimurium* CVD 1931 and *S. Enteritidis* CVD 1944. (A) OPA titers against wild-type *S. Typhimurium* D65 in serum samples from individual mice immunized with CVD 1931 (*S. Typhimurium* D65 Δ *guaBA* Δ *clpX*) or PBS. (B) OPA titers against wild-type *S. Enteritidis* S01 in serum samples from individual mice immunized with CVD 1944 (*S. Enteritidis* R11 Δ *guaBA* Δ *clpX*) or PBS. OPA titer was determined as the reciprocal of the highest serum dilution that produced >50% killing in comparison to bacteria incubated with BRC and HL-60 cells. Open circles, serum samples that were retested; *, $P < 0.05$ (Student's *t* test). Dashed line, the lowest dilution tested.




FIG 3 Correlation between anti-LPS serum IgG and OPA titers. Titers elicited by *S. Typhimurium* CVD 1931 (*S. Typhimurium* D65 Δ *guaBA* Δ *clpX*) (A) and *S. Enteritidis* CVD 1944 (*S. Enteritidis* R11 Δ *guaBA* Δ *clpX*) (B). Data were analyzed using Spearman's correlation coefficient.

centration of 12.5%, BRC was unable to mediate killing, as shown previously (16). When HL-60 cells were incubated with bacteria and serum without BRC, limited OPA titers were observed for immune mice (1:400) but not for nonimmune mice. However, when BRC was included in the assay, the OPA titer for immune sera increased >100-fold. It was therefore determined that a sublethal concentration of 12.5% BRC is required for optimal OPA in this assay.

Using our optimized OPA assay, we determined OPA titers of serum samples from individual mice immunized with live-attenuated NTS vaccines. We tested samples from groups that received the live-attenuated *S. Typhimurium* vaccine strain CVD 1931 (*S. Typhimurium* D65 Δ *guaBA* Δ *clpX*) against the target strain *S. Typhimurium* D65 from which the vaccine was derived. As expected, significantly higher OPA titers were measured in postvaccination sera than in samples from PBS-treated controls (Fig. 2A). Similarly, when we tested serum samples from mice immunized with the live-attenuated *S. Enteritidis* vaccine strain CVD 1944 (*S. Enteritidis* R11 Δ *guaBA* Δ *clpX*) against the virulent target strain *S. Enteritidis* S01, also isolated from the blood of a Malian child, we observed significantly higher OPA titers than for samples from mice treated with PBS (Fig. 2B). To account for possible lot-to-lot variation in BRC from the supplier, we picked 10 different serum samples from mice immunized with the live-attenuated *S. Enteritidis* vaccine strain CVD 1944 (*S. Enteritidis* R11 Δ *guaBA* Δ *clpX*) that were previously tested (Fig. 2B, open circles) for OPA titers with BRC lot 31325 and retested them in an OPA assay with BRC

TABLE 1 Seroconversion and anti-LPS serum IgG and OPA geometric mean titers (GMT)

Vaccine	Anti-LPS serum IgG titer			OPA titer		
	Preimmune GMT (95% CI)	Immune GMT (95% CI)	Seroconversion ^a	Preimmune GMT (95% CI)	Immune GMT (95% CI)	Seroconversion ^a
S. Typhimurium CVD 1931	12.5 (12.5–12.5)	4,337 (1,423–13,218)	20/21	100 (100–100)	71,223 (43,412–116,849)	21/21
S. Enteritidis CVD 1944	12.5 (12.5–12.5)	3,210 (1,104–9,329)	19/20	100 (100–100)	65,258 (36,925–115,331)	20/20

^a No. of mice with ≥ 4 -fold rise in titer.

lots 31325, 34636, and 19837. In our hands, we observed the OPA titers of the individual serum samples to be exactly the same for all three BRC lots. Furthermore, no differences in OPA titers were observed when the 10 immune serum samples were tested a second time on a different day with the same lot of BRC (lot 31325). A significant correlation was seen between OPA titers and anti-lipopolysaccharide (anti-LPS) serum IgG titers for *S. Typhimurium* and *S. Enteritidis* (Spearman's correlation coefficients of 0.85 and 0.88, respectively; $P < 0.0001$ for both) (Fig. 3). Using seroconversion as the parameter, all of the *S. Typhimurium*- and *S. Enteritidis*-vaccinated mice showed a >4 -fold increase in OPA titer compared to preimmune serum, while 20 of 21 *S. Typhimurium*-vaccinated mice and 19 of 20 *S. Enteritidis*-vaccinated mice seroconverted serum IgG anti-LPS titer (Table 1). The fact that the *S. Typhimurium*- and *S. Enteritidis*-vaccinated mice were protected (86% and 76% vaccine efficacies, respectively) against a lethal challenge (7) suggests that the OPA assay not only provides a measure of biological function but appears to be highly sensitive in detecting seroconversion following vaccination. The specificity of the OPA antibodies remains to be determined.

Robust, reproducible assays to measure immune responses elicited by candidate vaccines are required by regulatory agencies for eventual licensure. Here, we have adapted a well-characterized pneumococcus OPA assay accepted by regulatory agencies to evaluate pneumococcal conjugate vaccine formulations for measurement of OPA antibodies against *Salmonella*. Although the contribution of opsonophagocytic antibody activity in controlling *Salmonella* infections is unknown, the assay described herein may be used to characterize NTS vaccine immune responses, allowing quantification of functional antibody titers. The robustness of this assay, a measure of its reproducibility, was determined by using different lots of BRC and testing the assay on different days where the sera were subjected to repeated freeze-thaw cycles. We found the assay to be reproducible, with no differences observed in the OPA titers of the immune sera. We previously observed a similar robustness for SBA titers using immune mouse serum samples. Although SBA assays for NTS have been described, the need for an OPA assay is still compelling, as immune sera may lack SBA activity but still possess OPA that represents a correlate of protection for vaccines. Together with the SBA assay that we described previously, these assays may be useful in determining NTS correlates of protection, thereby allowing vaccine developers to predict vaccine performance.

ACKNOWLEDGMENT

This work was supported by a grant from NIH/NIAID (2 U54 AI057168) (principal investigator, M.M.L.).

FUNDING INFORMATION

This work, including the efforts of Girish Ramachandran, Mary Adetunke Boyd, Jennifer MacSwords, Ellen Higginson, Raphael Simon, James E. Galen, Marcela F. Pasetti, Myron M. Levine, and Sharon Mei Tennant, was funded by NIH (2 U54 AI057168).

REFERENCES

1. Feasey NA, Dougan G, Kingsley RA, Heyderman RS, Gordon MA. 2012. Invasive non-typhoidal *Salmonella* disease: an emerging and neglected tropical disease in Africa. *Lancet* 379:2489–2499. [http://dx.doi.org/10.1016/S0140-6736\(11\)61752-2](http://dx.doi.org/10.1016/S0140-6736(11)61752-2).
2. Ault A, Tennant SM, Gorres JP, Eckhaus M, Sandler NG, Roque A, Livio S, Bao S, Foulds KE, Kao SF, Roederer M, Schmidlein P, Boyd MA, Pasetti MF, Douek DC, Estes JD, Nabel GJ, Levine MM, Rao SS. 2013. Safety and tolerability of a live oral *Salmonella typhimurium* vaccine candidate in SIV-infected nonhuman primates. *Vaccine* 31:5879–5888. <http://dx.doi.org/10.1016/j.vaccine.2013.09.041>.
3. MacLennan CA, Martin LB, Micoli F. 2014. Vaccines against invasive *Salmonella* disease: current status and future directions. *Hum Vaccin Immunother* 10:1478–1493. <http://dx.doi.org/10.4161/hv.29054>.
4. Simon R, Tennant SM, Wang JY, Schmidlein PJ, Lees A, Ernst RK, Pasetti MF, Galen JE, Levine MM. 2011. *Salmonella enterica* serovar Enteritidis core O polysaccharide conjugated to H₂g_m flagellin as a candidate vaccine for protection against invasive infection with *S. Enteritidis*. *Infect Immun* 79:4240–4249. <http://dx.doi.org/10.1128/IAI.05484-11>.
5. Simon R, Wang JY, Boyd MA, Tulapurkar ME, Ramachandran G, Tennant SM, Pasetti M, Galen JE, Levine MM. 2013. Sustained protection in mice immunized with fractional doses of *Salmonella* Enteritidis core and O polysaccharide-flagellin glycoconjugates. *PLoS One* 8:e64680. <http://dx.doi.org/10.1371/journal.pone.0064680>.
6. Stefanetti G, Rondini S, Lanzillo L, Saul A, MacLennan CA, Micoli F. 2014. Impact of conjugation chemistry on the immunogenicity of *S. Typhimurium* conjugate vaccines. *Vaccine* 32:6122–6129. <http://dx.doi.org/10.1016/j.vaccine.2014.08.056>.
7. Tennant SM, Wang JY, Galen JE, Simon R, Pasetti MF, Gat O, Levine MM. 2011. Engineering and preclinical evaluation of attenuated nontyphoidal *Salmonella* strains serving as live oral vaccines and as reagent strains. *Infect Immun* 79:4175–4185. <http://dx.doi.org/10.1128/IAI.05278-11>.
8. Trebicka E, Shanmugam NK, Mikhailova A, Alter G, Cherayil BJ. 2014. Effect of human immunodeficiency virus infection on plasma bactericidal activity against *Salmonella enterica* serovar Typhimurium. *Clin Vaccine Immunol* 21:1437–1442. <http://dx.doi.org/10.1128/VI.00501-14>.
9. Trebicka E, Jacob S, Pirzai W, Hurley BP, Cherayil BJ. 2013. Role of antilipopolysaccharide antibodies in serum bactericidal activity against *Salmonella enterica* serovar Typhimurium in healthy adults and children in the United States. *Clin Vaccine Immunol* 20:1491–1498. <http://dx.doi.org/10.1128/VI.00289-13>.
10. Wahid R, Zafar SJ, McArthur MA, Pasetti MF, Levine MM, Sztein MB. 2014. Live oral *Salmonella enterica* serovar Typhi vaccines Ty21a and CVD 909 induce opsonophagocytic functional antibodies in humans that cross-react with *S. Paratyphi A* and *S. Paratyphi B*. *Clin Vaccine Immunol* 21:427–434. <http://dx.doi.org/10.1128/VI.00786-13>.
11. Gat O, Galen JE, Tennant S, Simon R, Blackwelder WC, Silverman DJ, Pasetti MF, Levine MM. 2011. Cell-associated flagella enhance the protection conferred by mucosally-administered attenuated *Salmonella* Para-

- typhi A vaccines. PLoS Negl Trop Dis 5:e1373. <http://dx.doi.org/10.1371/journal.pntd.0001373>.
12. Lindow JC, Fimlaid KA, Bunn JY, Kirkpatrick BD. 2011. Antibodies in action: role of human opsonins in killing *Salmonella enterica* serovar Typhi. Infect Immun 79:3188–3194. <http://dx.doi.org/10.1128/IAI.05081-11>.
 13. Pulickal AS, Gautam S, Clutterbuck EA, Thorson S, Basynat B, Adhikari N, Makepeace K, Rijpkema S, Borrow R, Farrar JJ, Pollard AJ. 2009. Kinetics of the natural, humoral immune response to *Salmonella enterica* serovar Typhi in Kathmandu, Nepal. Clin Vaccine Immunol 16: 1413–1419. <http://dx.doi.org/10.1128/CVI.00245-09>.
 14. MacLennan CA, Gondwe EN, Msefula CL, Kingsley RA, Thomson NR, White SA, Goodall M, Pickard DJ, Graham SM, Dougan G, Hart CA, Molyneux ME, Drayson MT. 2008. The neglected role of antibody in protection against bacteremia caused by nontyphoidal strains of *Salmonella* in African children. J Clin Invest 118:1553–1562. <http://dx.doi.org/10.1172/JCI33998>.
 15. Gondwe EN, Molyneux ME, Goodall M, Graham SM, Mastroeni P, Drayson MT, MacLennan CA. 2010. Importance of antibody and complement for oxidative burst and killing of invasive nontyphoidal *Salmonella* by blood cells in Africans. Proc Natl Acad Sci U S A 107:3070–3075. <http://dx.doi.org/10.1073/pnas.0910497107>.
 16. Boyd MA, Tennant SM, Saague VA, Simon R, Muhsen K, Ramachandran G, Cross AS, Galen JE, Pasetti MF, Levine MM. 2014. Serum bactericidal assays to evaluate typhoidal and nontyphoidal *Salmonella* vaccines. Clin Vaccine Immunol 21:712–721. <http://dx.doi.org/10.1128/CVI.00115-14>.
 17. Romero-Steiner S, Frasch C, Concepcion N, Goldblatt D, Kayhty H, Vakevainen M, Laferriere C, Wauters D, Nahm MH, Schinsky MF, Plikaytis BD, Carlone GM. 2003. Multilaboratory evaluation of a viability assay for measurement of opsonophagocytic antibodies specific to the capsular polysaccharides of *Streptococcus pneumoniae*. Clin Diagn Lab Immunol 10:1019–1024. <http://dx.doi.org/10.1128/CDLI.10.6.1019-1024.2003>.
 18. Romero-Steiner S, Frasch CE, Carlone G, Fleck RA, Goldblatt D, Nahm MH. 2006. Use of opsonophagocytosis for serological evaluation of pneumococcal vaccines. Clin Vaccine Immunol 13:165–169. <http://dx.doi.org/10.1128/CVI.13.2.165-169.2006>.
 19. Tennant SM, Diallo S, Levy H, Livio S, Sow SO, Tapia M, Fields PI, Mikoleit M, Tamboura B, Kotloff KL, Nataro JP, Galen JE, Levine MM. 2010. Identification by PCR of non-typhoidal *Salmonella enterica* serovars associated with invasive infections among febrile patients in Mali. PLoS Negl Trop Dis 4:e621. <http://dx.doi.org/10.1371/journal.pntd.0000621>.