

# Role of Arachidonic Acid and Its Metabolites in the Priming of NADPH Oxidase in Human Polymorphonuclear Leukocytes by Peritoneal Dialysis Effluent

I. DANIELS,<sup>1\*</sup> M. A. LINDSAY,<sup>1</sup> C. I. C. KEANY,<sup>1</sup> R. P. BURDEN,<sup>2</sup> J. FLETCHER,<sup>1</sup> AND A. P. HAYNES<sup>1</sup>

*Medical Research Centre<sup>1</sup> and Department of Renal Medicine,<sup>2</sup> City Hospital, Nottingham, United Kingdom*

Received 26 January 1998/Returned for modification 10 March 1998/Accepted 18 May 1998

Peritoneal dialysis effluent (PDE) contains a low-molecular-weight solute that will activate and prime the NADPH oxidase of human neutrophils via a phospholipase A<sub>2</sub> (PLA<sub>2</sub>)-dependent mechanism. Since the products of PLA<sub>2</sub> are known to activate and prime the oxidase we have investigated their role in the dialysis effluent-mediated activation and priming of human neutrophils. NADPH oxidase activity of PDE-primed and -unprimed neutrophils was measured by lucigenin-enhanced chemiluminescence in the presence of known inhibitors of the arachidonic acid cascade. Incubation of neutrophils with the nonselective PLA<sub>2</sub> inhibitor quinacrine (0 to 100 μM) reduced oxidase activity in both primed and unprimed cells. Furthermore, primed cells were more sensitive to the action of quinacrine than were unprimed cells. We were unable to determine the relative roles of secretory PLA<sub>2</sub> (sPLA<sub>2</sub>) and cytosolic PLA<sub>2</sub> (cPLA<sub>2</sub>) since the selective sPLA<sub>2</sub> inhibitor scalaradial (0 to 100 μM) inhibited oxidase activity in both groups of cells by similar degrees, while the specific cPLA<sub>2</sub> inhibitor AACO-CF<sub>3</sub> (0 to 50 μM) failed to affect activity in either group. Inhibition of platelet-activating factor (PAF), cyclooxygenase, and 5-lipoxygenase-activating protein by hexanolamino-PAF (0 to 25 μM), flurbiprofen (0 to 25 μM), and MK886 (0 to 5 μM), respectively, had no effect upon oxidase activity. However, the direct inhibition of 5-lipoxygenase by caffeic acid or lipoxin A<sub>4</sub> resulted in a similar concentration-dependent attenuation of oxidase activity in both primed and unprimed cells. Leukotriene B<sub>4</sub> (LTB<sub>4</sub>) release from primed neutrophils was comparable to that from unprimed cells with the exception of phorbol myristate acetate-stimulated cells, which released fivefold more LTB<sub>4</sub> than control. Taken together, these results suggest that it is arachidonic acid per se, and not its metabolites, that is important in priming of the neutrophil NADPH oxidase by dialysis effluent.

Recurrent bacterial peritonitis is a major cause of morbidity in patients undergoing continuous ambulatory peritoneal dialysis (CAPD). Polymorphonuclear leukocytes (PMN) are the predominant defense cells mobilized during an episode of infection; however, the intracellular killing mechanisms of these cells are impaired in the presence of peritoneal dialysis effluent (PDE) (18, 19). Paradoxically, PDE augments the release of secondary granules from PMN (11), causes low-level activation of the NADPH oxidase, and primes the latter response to both soluble and particulate stimuli (10). Subsequent investigation into the PMN signalling pathways affected by PDE suggest that both priming and activation involve phospholipase A<sub>2</sub> (PLA<sub>2</sub>) (28). The main products of PLA<sub>2</sub> activation in PMN are arachidonic acid (AA) and, if phosphatidylcholine is the substrate, platelet-activating factor (PAF). AA may be further metabolized by 5-lipoxygenase (5-LO) to yield leukotriene B<sub>4</sub> (LTB<sub>4</sub>) or by cyclooxygenase (COX) to yield prostaglandin. This latter reaction is thought to be minimal in PMN. Both PAF and LTB<sub>4</sub> are well recognized as being potent inflammatory mediators, and both are capable of priming (13, 22) and activating the NADPH oxidase (1, 46). Furthermore, AA is able to directly activate oxidase in cell (17) and cell-free systems (44) and to prime the response to subsequent stimulation by fMLP (40). We have therefore investigated the roles of PAF and AA and its metabolites in the priming and activation of the NADPH oxidase by PDE.

\* Corresponding author. Mailing address: Medical Research Centre, City Hospital, Hucknall Rd., Nottingham NG5 1PB, United Kingdom. Phone: (0115) 9627650, ext. 46509. Fax: (0115) 9602140. E-mail: ian.daniels@pmfmr.c.nottingham.ac.uk.

## MATERIALS AND METHODS

**Reagents.** Lucigenin, bovine serum albumin (BSA; fraction V, low endotoxin), quinacrine (PLA<sub>2</sub> inhibitor), flurbiprofen (COX inhibitor), and caffeic acid (lipoxygenase inhibitor) were obtained from Sigma Chemical Company Ltd., Poole, Dorset, United Kingdom. [5,6,8,9,11,12,14,15-<sup>3</sup>H]AA was obtained from Amersham International plc, Buckingham, United Kingdom. Lipoxin A<sub>4</sub> (LTB<sub>4</sub> receptor antagonist), hexanolamino-PAF (PAF receptor antagonist), arachidonyl trifluoromethyl ketone (AACO-CF<sub>3</sub>) (cytosolic PLA<sub>2</sub> [cPLA<sub>2</sub>] inhibitor) and scalaradial (secretory PLA<sub>2</sub> [sPLA<sub>2</sub>] inhibitor) were obtained from Cascade Biochemical, Reading, Berkshire, United Kingdom. MK886 (5-LO-activating protein [FLAP] inhibitor) was a generous gift from A. W. Ford-Hutchingson, Merck, Frost Centre, Quebec, Canada.

**Preparation of PMN.** Human peripheral blood PMN were prepared by standard methods (4). Fresh venous blood was added to EDTA (dipotassium salt) to a final concentration of 3.5 mM. Erythrocytes were sedimented on dextran, and the leukocyte-rich plasma was further purified over a Ficoll gradient (lymphocyte separation medium; Flow Laboratories, Herts, United Kingdom). The PMN-rich pellet was subjected to hypotonic lysis to remove the remaining erythrocytes and was washed twice in phosphate-buffered saline (PBS; pH 7.4). Purification of cells by this method routinely gave preparations of >99% viability as assessed by trypan blue exclusion and of >97% purity as assessed by examination of stained cytosin preparations. Cells were counted in a hemocytometer and were suspended at a concentration of 10<sup>7</sup> ml<sup>-1</sup> for lucigenin-enhanced chemiluminescence or 10<sup>8</sup> ml<sup>-1</sup> for LTB<sub>4</sub> determination by enzyme-linked immunosorbent assay (ELISA) (R & D Systems, Abingdon, Oxon, United Kingdom).

**PDE.** Six samples of PDE (1.36%, wt/vol, glucose) (Dianeal; Baxter Travenol Inc., Chicago, Ill.) were obtained from patients receiving CAPD after an intraperitoneal dwell of 4 h. All patients had been established on CAPD for more than 2 months, were not suffering infection, and had not received antibiotic therapy over the preceding 4 weeks. PDE was stored at -70°C until required. Before use PDE was filtered through a 0.2-μm-pore-size membrane and the pH was adjusted to 7.4 by the addition of HEPES to give a final concentration of 20 mM. The concentrations of creatinine and urea in PDE were determined by autoanalysis. The concentrations of protein endotoxin, and tumor necrosis factor alpha (TNF-α) and interleukin 1β (IL-1β) were determined by the method of Lowry et al. (29), the limulus amoebocyte lysate assay (Sigma Diagnostic Kit, Sigma Chemical Company) and ELISA, respectively.

**Determination of inhibitor toxicity.** The toxicity of the inhibitors and vehicle (ethanol) upon PMN over a 30-min period was determined by ATP bioluminescence (7) and confirmed by trypan blue exclusion. The ability of antagonists to scavenge superoxide anions was determined by using a xanthine-xanthine oxidase cell-free system (9).

**Determination of AA release.** PMN ( $10^7 \text{ ml}^{-1}$ ) were incubated with  $1 \mu\text{Ci}$  of [ $^3\text{H}$ ]AA  $\text{ml}^{-1}$  at  $37^\circ\text{C}$  for 60 min in PBS (without  $\text{Ca}^{2+}$ )–0.1% (wt/vol) BSA. Cells were washed three times in PBS and were suspended at a concentration of  $4 \times 10^7 \text{ ml}^{-1}$ . Under these conditions the percentage of label incorporated into PMN was  $31.33\% \pm 6.67\%$  (mean  $\pm$  standard error of the mean [SEM],  $n = 6$ ). The reaction mixture (200  $\mu\text{l}$ ) contained PBS (pH 7.4), 1 mM  $\text{CaCl}_2$ , 0.7 mM  $\text{MgCl}_2$ , 0.1% (wt/vol) BSA, inhibitor and/or vehicle, and PMN to give a final concentration of  $4 \times 10^6 \text{ ml}^{-1}$  (corresponding to  $260,188 \pm 23,981 \text{ cpm}$  [ $n = 6$ ]). PDEs were used at a concentration of 50% (vol/vol). The mixture was preincubated at  $37^\circ\text{C}$  for 10 min before the reaction was initiated by the addition of  $1 \mu\text{M}$  fMLP. After 20 min the reaction was terminated by the addition of 500  $\mu\text{l}$  of ice-cold physiological saline, and PMN were sedimented by centrifugation ( $12,000 \times g$  for 15 s). The amounts of [ $^3\text{H}$ ]AA in the pellet and supernatant were determined by liquid scintillation counting.

**Determination of superoxide anion generation.** Superoxide generation by PMN was determined by lucigenin-enhanced chemiluminescence in a plate-reading luminometer (Lumiscan; Labsystems, Basinstoke, United Kingdom). The reaction mixture (200  $\mu\text{l}$ ) contained 25  $\mu\text{M}$  lucigenin, PBS (pH 7.4), 1 mM  $\text{CaCl}_2$ , 0.7 mM  $\text{MgCl}_2$ , 0.1% (wt/vol) BSA, inhibitor and/or vehicle, and PMN to give a final concentration of  $10^6 \text{ ml}^{-1}$ . PDEs were used at a concentration of 50% vol/vol. The mixture was preincubated at  $37^\circ\text{C}$  for 10 min before the reaction was initiated by the addition of stimulus. fMLP, phorbol myristate acetate (PMA), and preopsonized *Staphylococcus epidermidis* were added to give final concentrations of  $1 \mu\text{M}$ ,  $10 \text{ ng ml}^{-1}$ , and  $2 \times 10^7 \text{ ml}^{-1}$ , respectively. Superoxide anion formation was taken as the integral of superoxide dismutase-inhibitible light output over the initial 30 min of the reaction.

**Determination of  $\text{LTB}_4$  generation.** The generation of  $\text{LTB}_4$  by PMN was determined by ELISA (R&D Systems). The reaction mixture (250  $\mu\text{l}$ ) contained PBS (pH 7.4), 1 mM  $\text{CaCl}_2$ , 0.7 mM  $\text{MgCl}_2$ , 0.1% (wt/vol) BSA, 50% (vol/vol) PDE, and PMN to give a final concentration of  $10^7 \text{ ml}^{-1}$ . The mixture was preincubated at  $37^\circ\text{C}$  for 10 min before the reaction was initiated by the addition of stimulus. fMLP, PMA, and preopsonized *S. epidermidis* were added to give final concentrations of  $1 \mu\text{M}$ ,  $10 \text{ ng ml}^{-1}$ , and  $2 \times 10^7 \text{ ml}^{-1}$ , respectively. The reaction was terminated by the addition of 13  $\mu\text{l}$  of ice-cold citric acid (0.035M) to reduce the pH to 5.5. The reaction mixture was spun at  $400 \times g$  and  $4^\circ\text{C}$  for 10 min. The supernatants were removed and stored at  $-80^\circ\text{C}$  before the assay for  $\text{LTB}_4$ .

**Statistical analysis.** All data are given as means  $\pm$  SEMs. Analysis was performed using the Mann-Whitney U test calculated by the computer multifunction statistics library package NWStatpak (Northwest Analytical Inc., Portland, Oreg.).

## RESULTS

The biochemical profiles (means  $\pm$  SEMs) of the six dialysis effluents employed in this study were as follows: protein,  $3.18 \pm 0.37 \text{ mg ml}^{-1}$  (range, 2.25 to  $4.90 \text{ mg ml}^{-1}$ ); endotoxin,  $71.25 \pm 9.22 \text{ pg ml}^{-1}$  (range, 57 to  $114 \text{ pg ml}^{-1}$ ); urea,  $18.18 \pm 1.56 \text{ mM}$  (range, 14.0 to  $25.4 \text{ mM}$ ); and creatinine,  $900 \pm 70 \mu\text{M}$  (range, 707 to  $1290 \mu\text{M}$ ). Levels of  $\text{TNF-}\alpha$  and  $\text{IL-1}\beta$  were below the level of detection by ELISA.

**Effects of inhibitors upon NADPH oxidase activity.** When concentrations of ethanol in the reaction mixture exceeded 0.1% (vol/vol) (ca. 20 mM) a significant reduction in PMN chemiluminescence was observed (50% inhibitory concentration [ $\text{IC}_{50}$ ] = 0.5% [vol/vol]). This inhibitory effect of ethanol was not affected by the presence of PDE and was not due to cell death since the release of ATP from PMN was constant. In subsequent experiments the final concentration of ethanol (used as the vehicle for inhibitors) did not exceed 0.05% (vol/vol). The release of ATP from PMN when incubated with all concentrations of inhibitors was comparable to the release observed from cells in buffer, suggesting that at the concentrations routinely used these compounds did not result in cell death (data not shown). The oxygen radical scavenging ability of the inhibitors employed in this study was assessed in a xanthine-xanthine oxidase cell-free system. At the concentrations used these compounds did not affect measurement of cell-free-generated superoxide (data not shown).




FIG. 1. Effect of quinacrine upon AA release from fMLP ( $1 \mu\text{M}$ )-challenged PMN. PMN were incubated in buffer (open squares) or PDE (closed squares). Results are expressed as mean ( $\pm$  SEM) percent inhibitions relative to control (no quinacrine) ( $n = 6$ , six dialysis effluents upon PMN from six donors). Absolute control values (100%) were  $1.90\% \pm 0.22\%$  and  $4.80\% \pm 0.55\%$  total radioactivity incorporated for unprimed and primed cells, respectively.

**Phospholipase  $\text{A}_2$  inhibitors.** AA generation in human PMN occurs predominantly through the action of  $\text{PLA}_2$  upon membrane phospholipids.  $\text{PLA}_2$  acts at the *sn-2* position of the glycerol backbone of phospholipids to yield free fatty acid (usually AA) and, if phosphatidylcholine is the substrate, *lysophosphatidylcholine*. Because of the pivotal role that this enzyme plays in AA generation we investigated its role in PDE-mediated priming and activation of human PMN by the use of three structurally distinct  $\text{PLA}_2$  inhibitors. Quinacrine, a widely used nonselective  $\text{PLA}_2$  inhibitor (50), inhibited release of AA from fMLP-stimulated PMN with an  $\text{IC}_{50}$  of approximately  $15 \mu\text{M}$  (Fig. 1). The compound was a poor inhibitor of basal superoxide generation in unstimulated PMN ( $\text{IC}_{50} = 100 \mu\text{M}$ ) but was 10-fold more effective if PMN were first primed with PDE ( $\text{IC}_{50} = 10 \mu\text{M}$ ) (Fig. 2a). The  $\text{IC}_{50}$  of quinacrine for receptor-mediated stimuli (fMLP and *S. epidermidis*) was  $20 \mu\text{M}$ , and this was reduced to  $10 \mu\text{M}$  in PDE-primed cells (Fig. 2b). Quinacrine inhibited superoxide generation by PMA-stimulated cells with a potency which was similar to those of fMLP and *S. epidermidis*; however, priming of PMN with PDE did not affect this response (data not shown). These observations suggest a role for  $\text{PLA}_2$  in both the activation and priming of PMN by PDE. In an attempt to further determine the form of  $\text{PLA}_2$  involved we used scylaradial, which at low concentrations ( $0.07 \mu\text{M}$ ) is a selective inhibitor of the 14-kDa *sPLA}\_2* (31), and AACOCF<sub>3</sub>, a trifluoromethyl ketone analogue of AA that acts as a selective, slow, tight-binding inhibitor of the 85-kDa *cPLA}\_2* (48). Incubation of PMN with scylaradial (0 to  $100 \mu\text{M}$ ) resulted in a concentration-dependent inhibition of both AA release (Fig. 3) and superoxide generation in primed and unprimed cells ( $\text{IC}_{50}$ s of 8, 8, 40, and  $8 \mu\text{M}$  for basal, fMLP, PMA, and *S. epidermidis*, respectively) (Fig. 4). Conversely, AACOCF<sub>3</sub> (0 to  $50 \mu\text{M}$ ) did not inhibit AA release (Fig. 5) or superoxide generation (Fig. 6) by either primed or unprimed cells under all conditions tested. Indeed at concentrations between 1 and  $10 \mu\text{M}$  superoxide generation was slightly stimulated for all stimuli (Fig. 6).

**PAF inhibitor.** The release of AA from phosphatidylcholine by  $\text{PLA}_2$  is accompanied by the concomitant formation of




FIG. 2. Effects of quinacrine upon superoxide generation of unstimulated PMN (a) and PMN challenged with 1 μM fMLP (b). PMN were incubated in buffer (open circles) or PDE (closed circles). Results are expressed as mean (± SEM) percent inhibitions relative to control (no quinacrine) (n = 6, six dialysis effluents upon PMN from six donors). Asterisks indicate P values of ≤0.05 (Mann-Whitney U test). Absolute control values (100%) were 28.85 ± 9.40 relative light units (RLU) and 115 ± 23.57 RLU (a) and 1,107 ± 28 RLU and 3,618 ± 153 RLU (b) for unprimed and primed cells, respectively.

lyso-PAF, which is subsequently acetylated to generate the bioactive PAF molecule. Since PAF is generated when PMN are activated by many stimuli (36) and will both prime (22) and activate (1) PMN, we investigated its role in PDE-mediated priming with the use of the PAF receptor antagonist hexanolamino-PAF (42). Concentrations of hexanolamino-PAF from 0 to 25 μM had no effect upon either primed or unprimed superoxide anion generation in response to all stimuli tested (unstimulated, fMLP, PMA, and *S. epidermidis*) (data not shown).

**5-LO inhibitors.** 5-LO is the predominant route by which AA is metabolized in human PMN. Because of the importance of this enzyme in the AA cascade we chose to use three func-

tionally distinct inhibitors to investigate the role of LTB<sub>4</sub> in PDE-primed PMN. The indole-type inhibitor MK886 inhibits FLAP, preventing the translocation of 5-LO from the cytosol to its active sight within the membrane (43). Caffeic acid inhibits LTB<sub>4</sub> generation by inactivating the active sight of 5-LO (26), and lipoxin A<sub>4</sub> is a reported LTB<sub>4</sub> receptor antagonist (6). Incubation of PMN with MK886 (0 to 5 μM) had no effect upon the release of superoxide in response to any of the stimuli used (unstimulated, fMLP, PMA, and *S. epidermidis*) (data not shown). PMN incubated with caffeic acid (0 to 200 μM) or lipoxin A<sub>4</sub> (0 to 5 μM) showed a concentration-dependent




FIG. 3. Effect of scalaradial upon AA release from fMLP (1 μM)-challenged PMN. PMN were incubated in buffer (open squares) or PDE (closed squares). Results are expressed as mean (± SEM) percent inhibitions relative to control (no scalaradial) (n = 6, six dialysis effluents upon PMN from six donors). Absolute control values (100%) were 1.80% ± 0.21% and 3.42% ± 0.37% total radioactivity incorporated for unprimed and primed cells, respectively.




FIG. 4. Effect of scalaradial upon superoxide generation by fMLP (1 μM)-challenged PMN. PMN were incubated in buffer (open circles) or PDE (closed circles). Results are expressed as mean (± SEM) percent inhibitions relative to control (no scalaradial) (n = 6, six dialysis effluents upon PMN from six donors). Similar profiles were obtained from unstimulated PMN and PMN challenged with *S. epidermidis* (2 × 10<sup>7</sup> ml<sup>-1</sup>) and PMA (10 ng ml<sup>-1</sup>). Absolute control values (100%) were 503 ± 42 relative light units (RLU) and 2,241 ± 224 RLU for unprimed and primed cells, respectively.




FIG. 5. Effect of AACO-CF<sub>3</sub> upon AA release from fMLP (1 μM)-challenged PMN. PMN were incubated in buffer (open squares) or PDE (closed squares). Results are expressed as mean (± SEM) percent inhibitions relative to control (no AACO-CF<sub>3</sub>) ( $n = 6$ , six dialysis effluents upon PMN from six donors). Absolute control values (100%) were 2.20% ± 0.25% and 4.62% ± 0.19% total radioactivity incorporated for unprimed and primed cells, respectively.

decrease in superoxide generation in response to all stimuli. The IC<sub>50</sub>s of caffeic acid and lipoxin A<sub>4</sub> (100 and 3 μM, respectively) were independent of the stimulus employed. Priming PMN with PDE prior to treatment with MK 886, caffeic acid, or lipoxin A<sub>4</sub> did not affect the results obtained.

**COX inhibitor.** COX is a ubiquitous mammalian enzyme that in PMN demonstrates a  $K_m$  for AA which is similar to that of lipoygenase. Activation of COX in PMN results in the formation of prostaglandin E<sub>2</sub> and thromboxin B<sub>2</sub>. We investigated the role of this enzyme in PDE-mediated priming and




FIG. 6. Effect of AACO-CF<sub>3</sub> upon superoxide generation from fMLP (1 μM)-challenged PMN. PMN were incubated in buffer (open circles) or PDE (closed circles). Results are expressed as mean (± SEM) percent inhibitions relative to (no AACO-CF<sub>3</sub>) ( $n = 6$ , six dialysis effluents upon PMN from six donors). Similar profiles were obtained from unstimulated PMN and PMN challenged with *S. epidermidis* ( $2 \times 10^7$  ml<sup>-1</sup>) and PMA (10 ng ml<sup>-1</sup>). Absolute control values (100%) were 747 ± 83 relative light units (RLU) and 1,842 ± 90 RLU for unprimed and primed cells, respectively.




FIG. 7. Generation of LTB<sub>4</sub> by unstimulated PMN and PMN challenged with fMLP (1 μM), *S. epidermidis* ( $2 \times 10^7$  ml<sup>-1</sup>), and PMA (10 ng ml<sup>-1</sup>). PMN were incubated in buffer (open bars) or PDE (hatched bars). Results are expressed as mean (± SEM) amount of LTB<sub>4</sub> released (ng ml<sup>-1</sup>) ( $n = 6$ , six dialysis effluents upon PMN from six donors). The asterisk indicates a  $P$  value of ≤ 0.05 (Mann-Whitney U test).

activation of human PMN with the COX inhibitor flurbiprofen. The generation of superoxide by PMN when unstimulated or stimulated by fMLP, PMA, or *S. epidermidis* was unaffected by the incubation of cells with the antagonist (0 to 25 μM) (data not shown). Furthermore, priming of PMN by PDE prior to treatment with the inhibitor did not affect the results (data not shown).

**ELISA measurement of LTB<sub>4</sub>.** In view of the important role of LTB<sub>4</sub> in mediating inflammatory responses, and with the consideration that we achieved various effects upon the generation of superoxide by using three different 5-LO antagonists, we directly quantified LTB<sub>4</sub> release from PMN by ELISA. The presence of PDE did not affect the measurement of known amounts of standard LTB<sub>4</sub>. Furthermore, PDE contained no LTB<sub>4</sub> per se (data not shown). The levels of LTB<sub>4</sub> released from unstimulated PMN and PMN stimulated by fMLP, PMA, and *S. epidermidis* were 13.70 ± 0.63, 39.30 ± 9.24, 47.30 ± 2.69, and 24.30 ± 5.35 ng ml<sup>-1</sup>, respectively (Fig. 7). In the presence of PDE these values remained unchanged with the exception of PMN stimulated by PMA. Under these conditions LTB<sub>4</sub> release was increased fivefold to 230.60 ± 41.00 ng ml<sup>-1</sup>.

## DISCUSSION

We have previously reported that incubation of PMN with 4-h intraperitoneal dwell, pH-corrected PDE results in low-level activation of the NADPH oxidase and primes this response to receptor-mediated stimuli (10), possibly by an effect upon specific granule release (11). Studies of the signal transduction systems affected by PDE demonstrate a profound activation of PLA<sub>2</sub> leading to an increase in the release of AA (28). In human PMN AA can directly activate the NADPH oxidase (17) and cause release of specific granules (23). Furthermore PAF (resulting from the liberation of AA from phosphatidylcholine) and the AA metabolite LTB<sub>4</sub> are potent activators and primers of many PMN functions (1, 13, 22, 46). With this in mind we have investigated the role of AA and its

metabolites in the priming and activation of human PMN by PDE.

A number of studies have investigated the role of PLA<sub>2</sub> in the activation of the NADPH oxidase in human PMN. The fact that superoxide generation in response to many stimuli is reduced upon incubation of cells with PLA<sub>2</sub> inhibitors and is restored by the addition of exogenous AA (20, 34, 37) demonstrates the importance of PLA<sub>2</sub> in the activation and maintenance of the NADPH oxidase. Using both quinacrine and scalaradial we were able to confirm a role for PLA<sub>2</sub> in PMN oxidase activation. Quinacrine reduced both superoxide generation and AA release from PMN in response to all stimuli tested. These data are in agreement with those of others who have demonstrated inhibition of superoxide generation of PMN by quinacrine when challenged with fMLP (37), PMA (20), and immune complexes (37). Our observation that PMN incubated with PDE were more sensitive to quinacrine prompted us to attempt to identify the form of PLA<sub>2</sub> activated by PDE by the use of two selective PLA<sub>2</sub> inhibitors. Incubation of PMN with scalaradial resulted in a concentration-dependent decrease in superoxide generation and AA release; however, the degrees of inhibition were similar in both primed and unprimed cells. These data would suggest that sPLA<sub>2</sub> does play a role in the generation of superoxide; however, since we report no difference between the IC<sub>50</sub> for the primed and unprimed cells, its role in PDE-mediated priming is questionable. Mayer et al. (32) have reported inhibition of superoxide generation by scalaradial in PMA-stimulated rat alveolar macrophage. In a manner similar to that for PDE, the priming of these macrophage with lipopolysaccharide (LPS) did not affect their results. It is generally accepted that it is cPLA<sub>2</sub> that is important in the release of AA from PMN (14). We investigated the role of cPLA<sub>2</sub> by using the arachidonyl trifluoromethyl ketone AACO-CF<sub>3</sub>. AACO-CF<sub>3</sub> is a potent inhibitor of cPLA<sub>2</sub> in cell-free systems (48). However, in the present study, exposure of PMN to the antagonist failed to inhibit AA release and resulted in a slight stimulation of the NADPH oxidase. These observations may reflect metabolism of the compound by PMN; indeed, recent reports have suggested that AACO-CF<sub>3</sub> is reduced to its noninhibitory alcohol upon incubation with many cell types (39).

It is well documented that exogenous PAF can stimulate oxidase activity and prime the response to subsequent stimulation by fMLP (1, 13). More recently, PAF receptor antagonists have been shown to inhibit superoxide production (47). We were, however, unable to demonstrate an effect of the PAF receptor antagonist hexanolamino-PAF upon either primed or unprimed superoxide generation by PMN when challenged with a variety of stimuli. The IC<sub>50</sub> of hexanolamino-PAF for superoxide generation in human macrophage is quoted at 52 μM (42). In this study, using ATP bioluminescence as an indicator of cell viability, we observed significant PMN death at concentrations of the antagonist exceeding 25 μM (data not shown). Our results suggest that PAF plays only a limited role in the generation of superoxide anion under the conditions employed in our studies. However, fMLP-stimulated PMN are known to generate PAF (36), and most of this PAF appears to remain within the cell (35). Since PAF may have an autocrine effect (33) and exert its effects by binding to the recently described high-affinity cytosolic PAF receptors (49), the efficiency of hexanolamino-PAF in blocking PAF-mediated responses may be compromised by its inability to penetrate the cell. The major pathway by which AA is metabolized in human PMN is via 5-LO, ultimately leading to the generation of LTB<sub>4</sub> (52). LTB<sub>4</sub> is released upon challenge of PMN with opsonized particles (5, 38), fMLP, and PMA (45). The inflammatory medi-

ator can directly activate aggregation, Ca<sup>2+</sup> mobilization, superoxide generation, and degranulation (46) and prime these responses to stimulation by agents such as fMLP (13). Because of its potential importance in mediating the inflammatory response we used three functionally distinct LTB<sub>4</sub> inhibitors to assess its role in PDE-induced priming and activation. In this study we obtained different results using these three inhibitors. Superoxide generation was inhibited with caffeic acid and lipoxin A<sub>4</sub> but not with MK886. These observations call into question the target selectivity of some of these antagonists. Recently, lipoxin A<sub>4</sub> has been shown to inhibit phosphoinositol hydrolysis in human PMN at concentrations similar to those employed in this study (16). The inhibition of superoxide generation that we report may therefore be a reflection of the inhibition of inositol-specific phospholipase C. No effect upon superoxide generation was seen with the FLAP inhibitor MK886. In agreement with others (43) we estimate the IC<sub>50</sub> of this compound for ionophore-stimulated LTB<sub>4</sub> release from human PMN to be approximately 120 nM (data not shown). However, we observed no reduction in superoxide generation even at concentrations of the antagonist (5 μM) that completely abolished LTB<sub>4</sub> release. We confirmed that LTB<sub>4</sub> plays little part in priming and activation of PMN by PDE by directly quantifying its release from cells by ELISA. In agreement with others we report that cells stimulated with fMLP, PMA, and *Staphylococcus* sp. (21, 30, 51) release only low levels of LTB<sub>4</sub> in comparison to the level released by ionophore-challenged cells. Only cells primed with PDE and subsequently stimulated by PMA release more LTB<sub>4</sub> than their relative controls. This observation may be a reflection of a combination of increased generation of AA (resulting from incubation of PMN with PDE) (28) and a direct activation of 5-LO by PMA. Several studies have reported that exposure of PMN to fresh (unused) dialysates results in a severe reduction in leukotriene release (24, 25). The data in our study confirm their suggestions that this effect is transient and probably the result of the low pH and high osmolality of fresh dialysates.

Flurbiprofen is well recognized as being a selective, potent inhibitor of both COX 1 and COX 11. In this study, concentrations of flurbiprofen of up to 25 μM had no effect upon oxidase activity of PMN when challenged with fMLP, PMA, or *S. epidermidis*. These observations are in keeping with those of others who report an ibuprofen IC<sub>50</sub> of 600 μM for fMLP-induced superoxide generation (37). These observations support the current thinking that although COX and 5-LO have similar K<sub>m</sub>s for AA, most if not all of the fatty acid generated during cell activation is metabolized by 5-LO (52).

The involvement of PLA<sub>2</sub> in primed stimulation is well documented. The enzyme appears to play a role in priming by diacyl- and alkylacylglycerol, inflammatory cytokines, and LPS (2, 3, 8, 12, 15, 27, 41). We were unable to detect appreciable concentrations of cytokines in the fluids employed in this study. Such an observation may not be considered unusual since the patients that we chose were not suffering infection at the time of dialysis. Our observation that only challenge with PMA results in elevated LTB<sub>4</sub> release from PDE-primed PMN is reminiscent of LPS priming (14, 15). However, the concentrations of endotoxin in the effluents used were below those required for priming (14), and unlike LPS, PDE-mediated priming of the oxidase is immediate with no requirement for incubation (10).

Taken together, we suggest that these results demonstrate that AA per se, and not its metabolites, is important in priming and activation of the NADPH oxidase by PDE. Further work to support a role for cPLA<sub>2</sub> in the generation of AA in PDE-primed human PMN and to study the chemical nature of the

molecule(s) responsible for these effects is currently under way in our laboratory.

#### ACKNOWLEDGMENTS

We thank the Trent Regional Health Authority, Sheffield, United Kingdom, and Baxters Healthcare Corp., Deerfield, Ill., for their support.

#### REFERENCES

- Bates, E. J., D. P. Harvey, and A. Ferrante. 1992. Inhibition of neutrophil respiratory burst and degranulation responses to platelet-activating factor by antagonists WEB 2086, CV 6209 and CV 3988. *Int. Arch. Allergy Immunol.* **97**:50-56.
- Bauldry, S. A., C. E. McCall, S. L. Cousart, and D. A. Bass. 1991. TNF $\alpha$  priming of PLA<sub>2</sub> activation in human neutrophils—an alternative mechanism of priming. *J. Immunol.* **149**:1277-1285.
- Bauldry, S. A., R. L. Wykle, and D. A. Bass. 1988. PLA<sub>2</sub> activation in human neutrophils. *J. Biol. Chem.* **263**:16787-16795.
- Boyum, A. 1968. Isolation of mononuclear cells and granulocytes from human blood. *Scand. J. Clin. Lab. Investig.* **21**(Suppl. 97):77-89.
- Classon, H.-E., U. Lundberg, and C. Malmsten. 1981. Serum coated zymosan stimulates the synthesis of LTB<sub>4</sub> in human PMN. Inhibition by cAMP. *Biochem. Biophys. Res. Commun.* **99**:1230-1237.
- Conti, P., M. Reale, R. C. Barbacane, M. R. Panara, and M. Bongrazio. 1991. Inhibition of LTB<sub>4</sub> in neutrophils by lipoxin A<sub>4</sub> and B<sub>4</sub>. *Agents Actions* **32**:85-87.
- Crouch, S. P. M., R. Kozlowski, K. J. Slater, and J. Fletcher. 1993. The use of ATP bioluminescence as a measure of cell proliferation and cytotoxicity. *J. Immunol. Methods* **160**:81-88.
- Dahinden, C. A., J. Zingg, F. E. Maly, and A. L. Weck. 1988. Leukotriene production in human neutrophils primed by recombinant human GMCSF and stimulated with the complement component C5a and fMLP as second signals. *J. Exp. Med.* **167**:1281-1293.
- Daniels, I., K. S. S. Bhatia, C. J. Porter, M. A. Lindsay, A. G. Morgan, R. P. Burden, and J. Fletcher. 1996. Hydrogen peroxide generation by polymorphonuclear leukocytes exposed to peritoneal dialysis effluent. *Clin. Diagn. Lab. Immunol.* **3**:682-688.
- Daniels, I., M. Lindsay, C. Porter, A. P. Haynes, J. Fletcher, and A. G. Morgan. 1993. Effect of peritoneal dialysis effluent on superoxide anion production by polymorphonuclear neutrophils. *Nephron* **64**:382-387.
- Daniels, I., S. P. M. Crouch, M. Lindsay, A. G. Morgan, R. P. Burden, and J. Fletcher. 1994. Primary and secondary granule release by polymorphonuclear leukocytes exposed to peritoneal dialysis effluent. *Clin. Diagn. Lab. Immunol.* **1**:227-231.
- Daniels, R. H., M. J. Finnen, M. E. Hill, and J. M. Lackie. 1992. Recombinant human monocyte IL8 primes NADPH-oxidase and PLA<sub>2</sub> activation in human neutrophils. *Immunology* **75**:157-163.
- Dewald, B., and M. Baggiolini. 1985. Activation of NADPH oxidase in human neutrophils. Synergism between fMLP and the neutrophil products PAF and LTB<sub>4</sub>. *Biochem. Biophys. Res. Commun.* **128**:297-304.
- Doerfler, M. E., J. Weiss, J. D. Clark, and P. Elsbach. 1994. Bacterial LPS primes human neutrophils for enhanced release of arachidonic acid and causes phosphorylation of an 85-kDa cytosolic PLA<sub>2</sub>. *J. Clin. Investig.* **93**:1583-1591.
- Doerfler, M. E., R. L. Danner, H. J. Shelhamer, and J. E. Parrillo. 1987. Bacterial LPS primes human neutrophils for enhanced production of LTB<sub>4</sub>. *J. Clin. Investig.* **83**:970-977.
- Grandordy, B. M., H. Lacroix, E. Mavoungou, S. Kriliso, A. E. G. Crea, B. W. Spur, and T. H. Lee. 1990. Lipoxin A<sub>4</sub> inhibits phosphoinositide hydrolysis in human neutrophils. *Biochem. Biophys. Res. Commun.* **167**:1022-1029.
- Hardy, S. J., A. Ferrante, A. Poulos, B. S. Robinson, D. W. Johnson, and A. W. Murray. 1994. Effect of exogenous fatty acids with greater than 22 carbon atoms (very long chain fatty acids) on superoxide production by human neutrophils. *J. Immunol.* **153**:1754-1761.
- Harvey, D. M., K. J. Sheppard, A. G. Morgan, and J. Fletcher. 1987. Effect of dialysis fluids on phagocytosis and killing by normal neutrophils. *J. Clin. Microbiol.* **25**:1424-1427.
- Harvey, D. M., K. J. Sheppard, A. G. Morgan, and J. Fletcher. 1988. Neutrophil function in patients on continuous ambulatory peritoneal dialysis. *Brit. J. Haematol.* **68**:273-278.
- Henderson, L. M., J. B. Chappell, and O. T. G. Jones. 1989. Superoxide generation is inhibited by PLA<sub>2</sub> inhibitors. Role for PLA<sub>2</sub> in the activation of the NADPH oxidase. *Biochem. J.* **264**:249-255.
- Henricks, P. A. J., M. E. Van Der Tol, F. Engels, F. P. Nijkamp, and J. Verhoef. 1986. Human polymorphonuclear leukocytes release LTB<sub>4</sub> during phagocytosis of *Staphylococcus aureus*. *Inflammation* **10**:37-47.
- Ingraham, L. M., T. D. Coates, J. M. Allen, C. P. Higgins, R. L. Baehner, and L. A. Boxer. 1982. Metabolic, membrane and functional responses of human PMN to PAF. *Blood* **59**:1259-1266.
- Jacobson, P. B., and D. J. Schrier. 1993. Regulation of CD11b/CD18 expression in human neutrophils by phospholipase A<sub>2</sub>. *J. Immunol.* **151**:5639-5652.
- Jorres, A., D. Jorres, N. Topley, G. M. Gahl, and A. Mahiout. 1991. Leukotriene release from peripheral and peritoneal leukocytes following exposure to peritoneal dialysis solutions. *Nephrol. Dial. Transplant* **6**:495-501.
- Jorres, A., D. Jorres, G. M. Gahl, M. Kessel, C. Muller, E. Kottgen, S. Serke, S. Schulz, and A. Mahiout. 1991. LTB<sub>4</sub> and TNF release from leukocytes: effect of peritoneal dialysis. *Nephron* **58**:276-282.
- Koshihara, Y., T. Neichi, S.-I. Murota, A.-N. Lao, Y. Fujimoto, and T. Tatsuno. 1984. Caffeic acid is a selective inhibitor for leukotriene biosynthesis. *Biochim. Biophys. Acta* **792**:92-97.
- Krump, E., and P. Borgeat. 1994. Kinetics of 5-LO activation, arachidonic acid release and leukotriene synthesis in human neutrophils: effects of GMCSF. *Biochem. Biophys. Acta* **1213**:135-139.
- Lindsay, M. A., I. Daniels, and J. Fletcher. 1997. Phospholipases and the activation and priming of neutrophils by peritoneal dialysis effluent. *Peritoneal Dial. Int.* **17**:471-479.
- Lowry, O. H., M. L. Rosenbrough, A. L. Farr, and R. J. Randall. 1951. Protein measurement with the Folin phenol reagent. *J. Biol. Chem.* **193**:265-275.
- Mahadevappa, V. G., and W. S. Powell. 1989. The metabolism of arachidonic and eicosapentaenoic acid in human neutrophils stimulated by A23187 and fMLP. *J. Cell. Biochem.* **40**:341-352.
- Marshall, L. A., J. D. Winkler, D. E. Griswold, B. Bolognese, A. Roshak, C.-M. Sung, E. F. Webb, and R. Jacobs. 1994. Effects of scalaradial, a type II PLA<sub>2</sub> inhibitor, on human neutrophil arachidonic acid mobilization and lipid mediator formation. *J. Pharmacol. Exp. Ther.* **268**:709-717.
- Mayer, A. M. S., S. Brenic, R. Stocker, and K. B. Glaser. 1995. Modulation of superoxide generation in in vivo lipopolysaccharide-primed rat alveolar macrophage by arachidonic acid and inhibitors of protein kinase C, PLA<sub>2</sub>, protein serine-threonine phosphatase(s), protein tyrosine kinase(s) and phosphatase(s). *J. Pharmacol. Exp. Ther.* **274**:427-436.
- McDonald, P. P., S. R. McColl, P. Braquet, and P. Borgeat. 1994. Autocrine enhancement of leukotriene synthesis by endogenous LTB<sub>4</sub> and platelet activating factor in human neutrophils. *Br. J. Pharmacol.* **111**:852-860.
- Muid, R. E., B. Twomey, and M. M. Dale. 1988. The effect of inhibition of both diacylglycerol metabolism and PLA<sub>2</sub> activity on superoxide generation by human neutrophils. *FEBS Lett.* **234**:235-240.
- Muller, S., and S. Nigam. 1992. Enhancement of staurosporine of PAF formation in fMLP challenged human PMN is mediated by intracellular PAF binding sites. *Biochem. Biophys. Res. Commun.* **189**:771-776.
- Muller, S., and S. Nigam. 1992. Arachidonic acid release and platelet-activating factor formation by staurosporine in human neutrophils challenged with n-formyl peptide. *Eur. J. Pharmacol.* **218**:251-258.
- Neal, T. M., M. C. M. Vissers, and C. C. W. Bouter. 1987. Inhibition by nonsteroidal anti-inflammatory drugs of superoxide production and granule enzyme release by polymorphonuclear leukocytes stimulated with immune complexes or fMLP. *Biochem. Pharmacol.* **36**:2511-2517.
- Palmer, R. M. J., and J. A. Salmon. 1983. Release of LTB<sub>4</sub> from human neutrophils and its relationship to degranulation induced by fMLP, serum treated zymosan and the ionophore A23187. *Immunology* **50**:65-73.
- Riendeau, D., J. Guay, P. K. Weech, F. Laliberte, J. Yergey, C. Li, S. Desmarais, H. Perrier, S. Liu, D. Nicol-Griiffith, and I. P. Street. 1994. Arachidonyl trifluoromethyl ketone, a potent inhibitor of 85-kDa PLA<sub>2</sub>, blocks production of arachidonic acid and 12-hydroxyeicosatetraenoic acid by calcium ionophore-challenged platelets. *J. Biol. Chem.* **269**:15619-15624.
- Roberts, P. J., S. L. Williams, and D. C. Linch. 1996. The regulation of neutrophil PLA<sub>2</sub> by granulocyte-macrophage colony-stimulating factor and its role in priming superoxide production. *Br. J. Haematol.* **92**:804-814.
- Rosenthal, M. D., and R. C. Franson. 1994. Separation of agonist-stimulated arachidonic mobilization from subsequent LTB<sub>4</sub> synthesis in human neutrophils: different effects of oleoylacylglycerol and PMA as priming agents. *J. Cell. Physiol.* **160**:522-530.
- Rouis, M., F. Nigon, and M. J. Chapman. 1988. Platelet activating factor is a potent stimulant of the production of active oxygen species by human monocyte-derived macrophages. *Biochem. Biophys. Res. Commun.* **156**:1293-1301.
- Rouzer, C. A., A. W. Ford-Hutchingson, H. E. Morton, and J. W. Gillard. 1990. MK886, a potent and specific leukotriene biosynthesis inhibitor blocks and reverses the membrane association of 5-lipoxygenase in ionophore-challenged leukocytes. *J. Biol. Chem.* **265**:1436-1442.
- Rubinek, T., and R. Levey. 1993. Arachidonic acid increases the activity of the assembled NADPH oxidase in cytoplasmic membranes and endosomes. *Biochim. Biophys. Acta* **1176**:51-58.
- Salari, H., P. Braquet, P. Naccache, and P. Borgeat. 1985. Characterization of effect of n-fMLP on leukotriene synthesis in human polymorphonuclear leukocytes. *Inflammation* **9**:127-138.
- Serhan, C. N., A. Radin, J. E. Smolen, H. Korchak, B. Samuelson, and G. Weissmann. 1982. LTB<sub>4</sub> is a complete secretagogue in human neutrophils: a kinetic analysis. *Biochem. Biophys. Res. Commun.* **107**:1006-1012.
- Steward, A. G., and T. Harris. 1991. Platelet-activating factor may partici-

- pate in signal transduction processes in rabbit leukocytes. *Lipids* **26**:1044–1049.
48. **Street, I. P., H.-K. Lin, F. Laliberte, F. Ghomashchi, Z. Wang, H. Perrier, N. M. Tremblay, Z. Huang, P. K. Weech, and M. H. Gelb.** 1993. Slow- and tight-binding inhibitors of the 85-kDa human PLA<sub>2</sub>. *Biochemistry* **32**:5935–5940.
49. **Svetlov, S., and S. Nigam.** 1993. Evidence for the presence of specific high affinity cytosolic binding sites for platelet activating factor in human neutrophils. *Biochim. Biophys. Acta* **190**:162–166.
50. **Vadas, P., and W. Pruzanski.** 1986. Biology of disease. Role of secretory PLA<sub>2</sub> in the pathobiology of disease. *Lab. Invest.* **55**:391–404.
51. **Waite, M., L. R. DeChatelet, L. King, and P. S. Shirley.** 1979. Phagocytosis-induced release of arachidonic acid from human neutrophils. *Biochem. Biophys. Res. Commun.* **90**:984–992.
52. **Walsh, C. E., B. M. Waite, M. J. Thomas, and L. R. DeChatelet.** 1981. Release and metabolism of arachidonic acid in human neutrophils. *J. Biol. Chem.* **256**:7228–7234.